



VIP2 rabbit pAb

Catalog No	BYab-11979
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	PPIP5K2 HISPPD1 KIAA0433 VIP2
Protein Name	VIP2
Immunogen	Synthesized peptide derived from human VIP2 AA range: 266-316
Specificity	This antibody detects endogenous levels of VIP2 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cytoplasm, cytosol .
Tissue Specificity	
Function	catalytic activity:ATP + 1D-myo-inositol 1,3,4,5,6-pentakisphosphate = ADP + diphospho-1D-myo-inositol tetrakisphosphate (isomeric configuration unknown).,catalytic activity:ATP + 1D-myo-inositol 5-diphosphate pentakisphosphate = ADP + 1D-myo-inositol bisdiphosphate tetrakisphosphate (isomeric configuration unknown).,catalytic activity:ATP + 1D-myo-inositol hexakisphosphate = ADP + 5-diphospho-1D-myo-inositol (1,2,3,4,6)pentakisphosphate.,caution:Although related to histidine acid phosphatases, it lacks the conserved active sites, suggesting that it has no phosphatase activity.,function:Bifunctional inositol kinase that catalyzes the formation of diphosphoinositol pentakisphosphate (InsP7 or PP-InsP5) and bi-diphosphoinositol tetrakisphosphate (InsP8 or PP2-InsP4). Converts inositolitol hexakisphosphate (InsP6) to InsP7. Also able to convert InsP7 to InsP8. Probably

Nanjing BYabscience technology Co.,Ltd



specifically mediates

Background

This gene encodes a member of the histidine acid phosphatase family of proteins. Despite containing a histidine acid phosphatase domain, the encoded protein functions as an inositol pyrophosphate kinase, and is thought to lack phosphatase activity. This kinase activity is the mechanism by which the encoded protein synthesizes high-energy inositol pyrophosphates, which act as signaling molecules that regulate cellular homeostasis and other processes. This gene may be associated with autism spectrum disorder in human patients. [provided by RefSeq, Sep 2016],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from THP-1 cells, primary antibody was diluted at 1:1000, 4° over night