



Cerebellin 1 Polyclonal Antibody

Catalog No	BYab-12698
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	CBLN1
Protein Name	Cerebellin-1
Immunogen	The antiserum was produced against synthesized peptide derived from human CBLN1. AA range:131-180
Specificity	Cerebellin 1 Polyclonal Antibody detects endogenous levels of Cerebellin 1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CBLN1; Cerebellin-1; Precerebellin
Observed Band	25kD
Cell Pathway	Secreted . Cell junction, synapse, postsynaptic cell membrane .
Tissue Specificity	In the Purkinje cells postsynaptic structures. In the cerebellum, cerebellin is much less abundant than [des-Ser1]-cerebellin.
Function	developmental stage:Low at birth, the cerebellin concentration increases between day 5 and 15, and reaches peak values between day 21 and 56.,function:Cerebellin exerts neuromodulatory functions. Directly stimulates norepinephrine release via the adenylate cyclase/PKA-dependent signaling pathway; and indirectly enhances adrenocortical secretion in vivo, through a paracrine mechanism involving medullary catecholamine release.,function:Cerebellin-1 is required for synapse integrity and synaptic plasticity. Required for the ER export and secretion of CBLN3.,PTM:The proteolytic processing to yield cerebellin seems to occur either prior to the secretion by presynaptic neurons and subsequent oligomerization or in some


Nanjing BYabscience technology Co.,Ltd




	other location after release of the mature protein.,similarity:Contains 1 C1q domain.,subcellular location:Might be bound to, or associated with, a membrane.,subunit:Homohexamer;
Background	This gene encodes a cerebellum-specific precursor protein, precerebellin, with similarity to the globular (non-collagen-like) domain of complement component C1qB. Precerebellin is processed to give rise to several derivatives, including the hexadecapeptide, cerebellin, which is highly enriched in postsynaptic structures of Purkinje cells. Cerebellin has also been found in human and rat adrenals, where it has been shown to enhance the secretory activity of this gland. [provided by RefSeq, Aug 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis of various cells using Cerebellin 1 Polyclonal Antibody



Western blot analysis of lysates from HUVEC cells, using CBLN1 Antibody. The lane on the right is blocked with the synthesized peptide.

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658