



TGFβ RIII Polyclonal Antibody

Catalog No	BYab-13697
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IF;ELISA
Gene Name	TGFBR3
Protein Name	Transforming growth factor beta receptor type 3
Immunogen	The antiserum was produced against synthesized peptide derived from human TGF beta Receptor III. AA range:801-850
Specificity	TGFβ RIII Polyclonal Antibody detects endogenous levels of TGFβ RIII protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/5000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TGFBR3; Transforming growth factor beta receptor type 3; TGF-beta receptor type 3; TGFR-3; Betaglycan; Transforming growth factor beta receptor III; TGF-beta receptor type III
Observed Band	
Cell Pathway	Secreted. Secreted, extracellular space. Cell membrane ; Single-pass type I membrane protein . Exists both as a membrane-bound form and as soluble form in serum and in the extracellular matrix.
Tissue Specificity	Blood,Brain,Placenta,Tongue,
Function	function:Binds to TGF-beta. Could be involved in capturing and retaining TGF-beta for presentation to the signaling receptors.,PTM:Extensively modified by glycosaminoglycan groups (GAG).,similarity:Contains 1 ZP domain.,subcellular location:Exists both as a membrane-bound form and as soluble form in serum and in the extracellular matrix.,subunit:Interacts with TCTEX1D4.,

Nanjing BYabscience technology Co.,Ltd



Background

This locus encodes the transforming growth factor (TGF)-beta type III receptor. The encoded receptor is a membrane proteoglycan that often functions as a co-receptor with other TGF-beta receptor superfamily members. Ectodomain shedding produces soluble TGFBR3, which may inhibit TGFB signaling. Decreased expression of this receptor has been observed in various cancers. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene.[provided by RefSeq, Sep 2010],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of NIH-3T3 cells using TGFβ RIII Polyclonal Antibody diluted at 1:1000



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using TGF beta Receptor III Antibody. The picture on the right is blocked with the synthesized peptide.