



INPP5J Polyclonal Antibody

Catalog No	BYab-14786
Isotype	IgG
Reactivity	Human;Rat
Applications	IHC;IF;ELISA
Gene Name	INPP5J
Protein Name	Phosphatidylinositol 4,5-bisphosphate 5-phosphatase A
Immunogen	Synthesized peptide derived from INPP5J . at AA range: 850-930
Specificity	INPP5J Polyclonal Antibody detects endogenous levels of INPP5J protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	INPP5J; PIB5PA; PIPP; Phosphatidylinositol 4; 5-bisphosphate 5-phosphatase A; Inositol polyphosphate 5-phosphatase J
Observed Band	
Cell Pathway	Cytoplasm . Predominantly localized to membrane ruffles. .
Tissue Specificity	Brain,Cerebellum,Heart,
Function	catalytic activity:1D-myo-inositol 1,3,4,5-tetrakisphosphate + H(2)O = 1D-myo-inositol 1,3,4-trisphosphate + phosphate.,catalytic activity:D-myo-inositol 1,4,5-trisphosphate + H(2)O = myo-inositol 1,4-bisphosphate + phosphate.,domain:The 5 Arg-Ser-Xaa-Ser-Xaa-Xaa (RSXSXX) motifs may constitute binding sites for the 14-3-3 protein.,function:Inositol 5-phosphatase, which converts inositol 1,4,5-trisphosphate to inositol 1,4-bisphosphate. Also converts phosphatidylinositol 4,5-bisphosphate to phosphatidylinositol 4-phosphate and inositol 1,3,4,5-tetrakisphosphate to inositol 1,3,4-trisphosphate in vitro. May be involved in modulation of the function of inositol and phosphatidylinositol polyphosphate-binding proteins that are present at membranes ruffles.,PTM:Phosphorylated at Ser/Thr residues.,similarity:Belongs

Nanjing BYabs science technology Co.,Ltd



to the inositol-1,4,5-trisphosphate 5-phosphatase type II family.,subcellular l

Background

catalytic activity:1D-myo-inositol 1,3,4,5-tetrakisphosphate + H(2)O = 1D-myo-inositol 1,3,4-trisphosphate + phosphate.,catalytic activity:D-myo-inositol 1,4,5-trisphosphate + H(2)O = myo-inositol 1,4-bisphosphate + phosphate.,domain:The 5 Arg-Ser-Xaa-Ser-Xaa-Xaa (RSXSXX) motifs may constitute binding sites for the 14-3-3 protein.,function:Inositol 5-phosphatase, which converts inositol 1,4,5-trisphosphate to inositol 1,4-bisphosphate. Also converts phosphatidylinositol 4,5-bisphosphate to phosphatidylinositol 4-phosphate and inositol 1,3,4,5-tetrakisphosphate to inositol 1,3,4-trisphosphate in vitro. May be involved in modulation of the function of inositol and phosphatidylinositol polyphosphate-binding proteins that are present at membranes ruffles.,PTM:Phosphorylated at Ser/Thr residues.,similarity:Belongs to the inositol-1,4,5-trisphosphate 5-phosphatase type II family.,subcellular location:Predominantly localized to membrane ruffles.,


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658