



HDAC6 (ABT-HDAC6) mouse mAb

Catalog No	BYab-15223
Isotype	IgG
Reactivity	Human; Predict react with Mouse
Applications	IHC;WB;IF
Gene Name	HDAC6 KIAA0901 JM21
Protein Name	Histone deacetylase 6 (HD6) (EC 3.5.1.98)
Immunogen	Synthesized peptide derived from human HDAC6
Specificity	This antibody detects endogenous levels of human HDAC6. Heat-induced epitope retrieval (HIER) Citrate buffer of pH6.0 was highly recommended as antigen repair method in paraffin section
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Mouse, Monoclonal/IgG1, Kappa
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1:100-500, WB 1:200-1000. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cytoplasm . Cytoplasm, cytoskeleton . Nucleus . Perikaryon . Cell projection, dendrite . Cell projection, axon . It is mainly cytoplasmic, where it is associated with microtubules. .
Tissue Specificity	Brain,Epithelium,Kidney,Muscle,Ovary,Placenta,
Function	catalytic activity:Hydrolysis of an N(6)-acetyl-lysine residue of a histone to yield a deacetylated histone.,function:Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes (By similarity). Plays a central role in microtubule-dependent cell motility via deacetylation of tubulin.,PTM:Sumoylated in vitro.,PTM:Ubiquitinated. Its polyubiquitination however does not lead to its degradation.,similarity:Belongs to


Nanjing BYabscience technology Co.,Ltd




	the histone deacetylase family. Type 2 subfamily.,similarity:Contains 1 UBP-type zinc finger.,subcellular location:It is mainly cytoplasmic, where it is associated with microtubules
Background	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It contains an internal duplication of two catalytic domains which appear to function independently of each other. This protein possesses histone deacetylase activity and represses transcription. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images




Human kidney tissue was stained with Anti-HDAC6 (ABT-HDAC6) Antibody



Human kidney tissue was stained with Anti-HDAC6 (ABT-HDAC6) Antibody



Human testis tissue was stained with Anti-HDAC6 (ABT-HDAC6) Antibody



Whole cell lysates were separated by 10% SDS-PAGE, and the membrane was blotted with anti-HDAC6 (ABT-HDAC6)antibody. The HRP-conjugated Goat anti-Mouse IgG(H + L) antibody was used to detect the antibody. Lane 1: HeLa Predicted band size: 131kDa Observed band size: 130,180kDa