



Cleaved-ITI-H4 70k (R661) Polyclonal Antibody


Catalog No	BYab-15817
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	IHC;IF;ELISA
Gene Name	ITIH4
Protein Name	Inter-alpha-trypsin inhibitor heavy chain H4
Immunogen	The antiserum was produced against synthesized peptide derived from human ITIH4. AA range:612-661
Specificity	Cleaved-ITI-H4 70k (R661) Polyclonal Antibody detects endogenous levels of fragment of activated ITI-H4 70k protein resulting from cleavage adjacent to R661.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/10000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ITIH4; IHRP; ITIHL1; PK120; Inter-alpha-trypsin inhibitor heavy chain H4; ITI heavy chain H4; ITI-HC4; Inter-alpha-inhibitor heavy chain 4; Inter-alpha-trypsin inhibitor family heavy chain-related protein; IHRP; Plasma kallikrein sensitive
Observed Band	
Cell Pathway	Secreted.
Tissue Specificity	Liver specific.
Function	function:May be involved in acute phase reactions.,induction:Levels increase significantly in the sera of patients after different surgical trauma.,PTM:Appears to be both N- and O-glycosylated.,PTM:Cleaved by plasma kallikrein to yield 100 kDa and 35 kDa fragments, and the resulting 100 kDa fragment is further converted to a 70 kDa fragment.,PTM:The O-linked polysaccharides on Ser-696, Thr-701 and Ser-702 are probably the mucin type linked to GalNAc.,similarity:Belongs to the ITIH family.,similarity:Contains 1 VWFA domain.,tissue specificity:Liver specific.,

Nanjing BYabscience technology Co.,Ltd



Background	The protein encoded by this gene is secreted into the blood, where it is cleaved by plasma kallikrein into two smaller forms. Expression of this gene has been detected only in liver, and it seems to be upregulated during surgical trauma. This gene is part of a cluster of similar genes on chromosome 3. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2009],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).