



Cyclin D3 Monoclonal Antibody

Catalog No	BYab-16556
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	CCND3
Protein Name	G1/S-specific cyclin-D3
Immunogen	Purified recombinant fragment of human Cyclin D3 expressed in E. Coli.
Specificity	Cyclin D3 Monoclonal Antibody detects endogenous levels of Cyclin D3 protein.
Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CCND3; G1/S-specific cyclin-D3
Observed Band	
Cell Pathway	Nucleus . Cytoplasm .
Tissue Specificity	Brain,Lung,Placenta,Synovial membrane tissue,Umbilical cord blood,
Function	function:Essential for the control of the cell cycle at the G1/S (start) transition. Potentiates the transcriptional activity of ATF5.,similarity:Belongs to the cyclin family.,similarity:Belongs to the cyclin family. Cyclin D subfamily.,subunit:Interacts with the CDK4 and CDK6 protein kinases to form a serine/threonine kinase holoenzyme complex. The cyclin subunit imparts substrate specificity to the complex. Interacts with ATF5. Interacts with EIF3K.,
Background	The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activitiy is required for

Nanjing BYabscience technology Co.,Ltd



cell cycle G1/S transition. This protein has been shown to interact with and be involved in the phosphorylation of tumor suppressor protein Rb. The CDK4 activity associated with this cyclin was reported to be necessary for cell cycle progression through G2 phase into mitosis after UV radiation. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2008],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis using Cyclin D3 Monoclonal Antibody against NIH/3T3 (1) and Jurkat (2) cell lysate.



Nanjing BYabs science technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658