



Contactin 5 Polyclonal Antibody

Catalog No	BYab-17012
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	IF;IHC;ELISA
Gene Name	CNTN5
Protein Name	Contactin-5
Immunogen	The antiserum was produced against synthesized peptide derived from human CNTN5. AA range:891-940
Specificity	Contactin 5 Polyclonal Antibody detects endogenous levels of Contactin 5 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CNTN5; Contactin-5; Neural recognition molecule NB-2; hNB-2
Observed Band	
Cell Pathway	Cell membrane ; Lipid-anchor, GPI-anchor .
Tissue Specificity	Expressed in brain and kidney and at very low level in placenta. Not expressed in other tissues. In brain, it is highly expressed in the occipital lobe, amygdala, cerebral cortex, frontal lobe, thalamus and temporal lobe. Expressed at moderate level in the cerebellum, substantia nigra, putamen, medulla and hippocampus. Weakly expressed in the spinal cord and caudate nucleus. Weakly or not expressed in the corpus callosum.
Function	function:Contactins mediate cell surface interactions during nervous system development. Has some neurite outgrowth-promoting activity in the cerebral cortical neurons but not in hippocampal neurons. Probably involved in neuronal activity in the auditory system.,similarity:Belongs to the immunoglobulin superfamily. Contactin family.,similarity:Contains 4 fibronectin type-III domains.,similarity:Contains 6 Ig-like C2-type (immunoglobulin-like) domains.,tissue specificity:Expressed in brain and kidney and at very low level in

Nanjing BYabscience technology Co.,Ltd



placenta. Not expressed in other tissues. In brain, it is highly expressed in the occipital lobe, amygdala, cerebral cortex, frontal lobe, thalamus and temporal lobe. Expressed at moderate level in the cerebellum, substantia nigra, putamen, medulla and hippocampus. Weakly expressed in the spinal cord and caudate nucleus. Weakly or not expressed in the corpus callosum.

Background

The protein encoded by this gene is a member of the immunoglobulin superfamily, and contactin family, which mediate cell surface interactions during nervous system development. This protein is a glycosylphosphatidylinositol (GPI)-anchored neuronal membrane protein that functions as a cell adhesion molecule. It may play a role in the formation of axon connections in the developing nervous system. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Aug 2011],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunofluorescence analysis of A549 cells, using CNTN5 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658