



ZIP8 rabbit pAb

货号:	BYab-17710
同型:	IgG
靶点:	SLC39A8
基因名称:	SLC39A8 BIGM103 ZIP8 PP3105
蛋白名称:	Zinc transporter ZIP8 (BCG-induced integral membrane protein in monocyte clone 103 protein) (LIV-1 subfamily of ZIP zinc transporter 6) (LZT-Hs6) (Solute carrier family 39 member 8) (Zrt- and Irt-like
免疫原:	Synthesized peptide derived from human ZIP8
特异性:	This antibody detects endogenous levels of ZIP8 at Human, Mouse, Rat
组成:	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
来源:	Polyclonal, Rabbit, IgG
稀释:	WB 1:500-2000
纯化工艺:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
浓度:	1 mg/ml
储存:	-15°C to -25°C/1 year (Do not lower than -25°C)
分子量:	51kD
功能:	Electroneutral transporter of the plasma membrane mediating the cellular uptake of zinc and manganese, two divalent metal cations important for development, tissue homeostasis or immunity. Functions as an energy-dependent symporter, transporting through the membranes an electroneutral complex composed of a divalent metal cation, a bicarbonate and a selenite anion or yet a metal cation and two bicarbonate anions. May also transport iron, mercury and cobalt through membranes. Beside these endogenous cellular substrates, also imports cadmium a non-essential metal which is cytotoxic and carcinogenic. Through zinc import, indirectly regulates the metal-dependent transcription factor MTF1 and the expression of some metalloproteases involved in cartilage catabolism and also probably heart development. Also indirectly regulates the expression of proteins involved in cell morphology and cyt
细胞定位:	Cell membrane; Multi-pass membrane protein. Lysosome membrane; Multi-pass membrane protein. Apical cell membrane; Multi-pass membrane protein. Basolateral cell membrane; Multi-pass membrane protein. Localizes to the lysosome of activated T-cells (PubMed:19401385). A large fraction of the protein is found intracellularly in microvascular capillary endothelial cells that constitute the blood-brain barrier (PubMed:31699897). Localized and functional at both apical and basolateral membranes of microvascular capillary endothelial cells that constitute the blood-brain barrier (PubMed:31699897).
组织表达:	Ubiquitously expressed (PubMed:12504855, PubMed:22898811, PubMed:28056086, PubMed:31699897). Expressed in thymus, placenta, lung,

Nanjing BYabs science technology Co., Ltd



liver, pancreas, salivary gland and, to a lower extent, in spleen, testis, ovary, small intestine, colon, leukocyte, heart. Highest expression is observed in pancreas (PubMed:12504855). Expressed by macrophages (at protein level) (PubMed:28056086). Expressed by microvascular capillary endothelial cells that constitute the blood-brain barrier (at protein level) (PubMed:31699897).

有关注意事项:

Avoid repeated freezing and thawing!

使用建议:

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images