



ABCE1 rabbit pAb

货号	BYab-17937
同位型	IgG
应用	WB
种属	Human;Mouse
靶点	ABCE1
基因名称	ABCE1 RLI RNASEL1 RNASELI RNS4I OK/SW-cl.40
蛋白名称	ATP-binding cassette sub-family E member 1 (2'-5'-oligoadenylate-binding protein) (HuHP68) (RNase L inhibitor) (Ribonuclease 4 inhibitor) (RNS4I)
免疫原	Synthesized peptide derived from human ABCE1
特异性	This antibody detects endogenous levels of ABCE1 at Human, Mouse
组成	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
来源	Polyclonal, Rabbit,IgG
稀释	WB 1:500-2000
纯化工艺	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
分子量	66kD
功能	Cotranslational quality control factor involved in the No-Go Decay (NGD) pathway . Together with PELO and HBS1L, is required for 48S complex formation from 80S ribosomes and dissociation of vacant 80S ribosomes . Together with PELO and HBS1L, recognizes stalled ribosomes and promotes dissociation of elongation complexes assembled on non-stop mRNAs; this triggers endonucleolytic cleavage of the mRNA, a mechanism to release non-functional ribosomes and to degrade damaged mRNAs as part of the No-Go Decay (NGD) pathway . Plays a role in the regulation of mRNA turnover (By similarity). Plays a role in quality control of translation of mitochondrial outer membrane-localized mRNA . As part of the PINK1-regulated signaling, ubiquitinated by CNOT4 upon mitochondria damage; this modification generates polyubiquitin signals that recruit autophagy receptors to the mitochondrial outer membrane and i
细胞定位	Cytoplasm . Mitochondrion . Localized to clusters of virus formation at the plasma membrane.
浓度	1 mg/ml
储存	-15°C to -25°C/1 year(Do not lower than -25°C)
有关注意事项	Avoid repeated freezing and thawing!

Nanjing BYabs science technology Co.,Ltd



使用建议

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images