



MGAT3 rabbit pAb

货号	BYab-17987
同位型	IgG
应用	WB
种属	Human;Rat
靶点	MGAT3
基因名称	MGAT3 GGNT3
蛋白名称	Beta-1,4-mannosyl-glycoprotein 4-beta-N-acetylglucosaminyltransferase (EC 2.4.1.144) (N-glycosyl-oligosaccharide-glycoprotein N-acetylglucosaminyltransferase III) (GNT-III) (GlcNAc-T III) (N-acetylglu
免疫原	Synthesized peptide derived from human MGAT3
特异性	This antibody detects endogenous levels of MGAT3 at Human,Rat
组成	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
来源	Polyclonal, Rabbit,IgG
稀释	WB 1:500-2000
纯化工艺	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
分子量	59kD
功能	It is involved in the regulation of the biosynthesis and biological function of glycoprotein oligosaccharides. Catalyzes the addition of N-acetylglucosamine in beta 1-4 linkage to the beta-linked mannose of the trimannosyl core of N-linked sugar chains, called bisecting N-acetylglucosamine (GlcNAc). It is one of the most important enzymes involved in the regulation of the biosynthesis of glycoprotein oligosaccharides. The addition of this bisecting GlcNAc residue alters not only the composition, but also the conformation of the N-glycan. The introduction of the bisecting GlcNAc residue results in the suppression of further processing and elongation of N-glycans, precluding the formation of beta-1,6 GlcNAc branching, catalyzed by MGAT5 since it is unable to use the bisected oligosaccharide as a substrate . Addition of bisecting N-acetylglucosamine to CDH1/E-cadherin modulates CDH1 cell m
细胞定位	Golgi apparatus membrane; Single-pass type II membrane protein .
浓度	1 mg/ml
储存	-15°C to -25°C/1 year(Do not lower than -25°C)
有关注意事项	Avoid repeated freezing and thawing!

Nanjing BYabs science technology Co.,Ltd



使用建议

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images