



BCSC-1 Monoclonal Antibody

Catalog No	BYmab-00326
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	VWA5A
Protein Name	von Willebrand factor A domain-containing protein 5A
Immunogen	The antiserum was produced against synthesized peptide derived from human LHR2A. AA range:371-420
Specificity	BCSC-1 Monoclonal Antibody detects endogenous levels of BCSC-1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	VWA5A; BCSC1; LOH11CR2A; von Willebrand factor A domain-containing protein 5A; Breast cancer suppressor candidate 1; BCSC-1; Loss of heterozygosity 11 chromosomal region 2 gene A protein
Observed Band	86kD
Cell Pathway	nucleus,nucleoplasm,
Tissue Specificity	Expressed at low level in many tissues. Not expressed in 80% of tumor cell lines tested.
Function	
Background	VWA5A (von Willebrand factor A domain containing 5A), also known as BCSC-1 (breast cancer suppressor candidate 1) or LOH11CR2A (loss of heterozygosity 11 chromosomal region 2 gene A protein), is a 786 amino acid protein containing one VIT domain and one VWFA domain. VWA5A is expressed at low levels in various tissues, with no expression found in 80% of tumor cell lines. Likely acting as a tumor suppressor gene, deletion of VWA5A leads to loss of heterozygosity (LOH)

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



in breast and ovarian tumors, and may have an important role as a potential gene
therapy target. Abnormal expression of VWA5A may lead to an increase in
adhesion of CNE-2L2 cells associated with an increase in expression of
E-cadherin, alpha-catenin, and p53, resulting in a decrease of malignant activity in
cells with ectopic expression of VWA5A. Existing as four alternatively spliced
isoforms, the gene encoding VWA5A maps to human chromosome 11q24.2.


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using BCSC-1 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd