



## BTF Monoclonal Antibody

Catalog No         BYmab-00338           Isotype         IgG           Reactivity         Human;Mouse;Rat           Applications         WB           Gene Name         BCLAF1           Protein Name         Bcl-2-associated transcription factor 1           Immunogen         Breatiserum was produced against synthesized peptide derived from human BCLAF1. AA range:281-330           Specificity         BTF Monoclonal Antibody detects endogenous levels of BTF protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         290%           Storage Stability         -20°C/1 year           Synonyms         BCLAF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf           Observed Band         106kD           Cell Pathway         Cytoplasm. Nucleus. Nucleus speckle . Nucleus, nucleoplasm .           Tissue Specificity         Ubiquitous.           Function         function:Death-promoting transcriptional repressor. sequence caution:Contaminating sequence. Potential poly-A sequence, sub		
Reactivity Applications WB Gene Name BCLAF1 Protein Name Bcl-2-associated transcription factor 1 Immunogen The antiserum was produced against synthesized peptide derived from human BCLAF1. AA range:281-330 Specificity BTF Monoclonal Antibody detects endogenous levels of BTF protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms BCLAF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf Observed Band 106kD Cell Pathway Cytoplasm. Nucleus. Nucleus speckle. Nucleus, nucleoplasm. Tissue Specificity Ubiquitous. Function function:Death-promoting transcriptional repressor, sequence caution:Contaminating sequence. Potential poly-A sequence, subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA, tissue specificity. Ubiquitous. This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redstributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Catalog No	BYmab-00338
Applications  Gene Name  BCLAF1  Protein Name  Bcl-2-associated transcription factor 1  Immunogen  The antiserum was produced against synthesized peptide derived from human BCLAF1. AA range:281-330  Specificity  BTF Monoclonal Antibody detects endogenous levels of BTF protein.  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  290%  Storage Stability  -20°C/1 year  Synonyms  BCLAF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf  Observed Band  106kD  Cell Pathway  Cytoplasm. Nucleus. Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity  Ubiquitous.  Function  function-Death-promoting transcriptional repressor, sequence caution: Contaminating sequence. Potential poly-A sequence, subunit: Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA, tissue specificity. Ubiquitous.  Background  This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apportosis. Multiple	Isotype	IgG
Gene Name Protein Name BcLaF1 Bcl-2-associated transcription factor 1  Immunogen The antiserum was produced against synthesized peptide derived from human BcLaF1. AA range:281-330 Specificity BTF Monoclonal Antibody detects endogenous levels of BTF protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse,lgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms BcLaF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf  Observed Band 106kD Cell Pathway Cytoplasm. Nucleus. Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity Ubiquitous.  Function function:Death-promoting transcriptional repressor.,sequence caution:Contaminating sequence. Potential poly-A sequence, subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA, tissue specificity:Ubiquitous.  Background This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergroing apoptosis. Multiples	Reactivity	Human;Mouse;Rat
Protein Name Bcl-2-associated transcription factor 1 Immunogen The antiserum was produced against synthesized peptide derived from human BCLAF1. AA range:281-330 Specificity BTF Monoclonal Antibody detects endogenous levels of BTF protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms BCLAF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf Observed Band 106kD Cell Pathway Cytoplasm. Nucleus. Nucleus speckle . Nucleus, nucleoplasm . Tissue Specificity Ubiquitous. Function function: Death-promoting transcriptional repressor, sequence caution: Contaminating sequence. Potential poly-A sequence, subunit: Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA, tissue specificity; Ubiquitous. This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of this protein induces apoptosis, which can be suppressed by co-expression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Applications	WB
Immunogen  The antiserum was produced against synthesized peptide derived from human BCLAF1. AA range:281-330  Specificity  BTF Monoclonal Antibody detects endogenous levels of BTF protein.  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source  Monoclonal, Mouse, lgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  BCLAF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf  Observed Band  106kD  Cell Pathway  Cytoplasm. Nucleus. Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity  Ubiquitous.  Function  function:Death-promoting transcriptional repressor, sequence caution:Contaminating sequence. Potential poly-A sequence, subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA.,tissue specificity. Ubiquitous.  Background  This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of this protein induces apoptosis, which can be suppressed by co-expression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Gene Name	BCLAF1
BCLAF1. AA range:281-330  Specificity BTF Monoclonal Antibody detects endogenous levels of BTF protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms BCLAF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf  Observed Band 106kD  Cell Pathway Cytoplasm. Nucleus. Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity Ubiquitous.  Function function: Death-promoting transcriptional repressor .sequence caution: Contaminating sequence. Potential poly-A sequence. subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA, tissue specificity: Ubiquitous.  Background This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of BCL2 proteins. The protein localizes to dot-like structures throughout apoptosis, Miltiple to a zone near the nuclear envelope in cells undergone apoptosis, Multiple	Protein Name	Bcl-2-associated transcription factor 1
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms BCLAF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf  Observed Band 106kD  Cell Pathway Cytoplasm. Nucleus. Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity Ubiquitous.  Function function:Death-promoting transcriptional repressor.,sequence caution:Contaminating sequence. Potential poly-A sequence, subunit:Interacts with Bcl-2 related profeins, EMD, with the adenovirus E1B 19 kDa protein and with DNA., tissue specificity:Ubiquitous.,  Background This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of Btl2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Immunogen	
Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         BCLAF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf           Observed Band         106kD           Cell Pathway         Cytoplasm. Nucleus. Nucleus speckle . Nucleus, nucleoplasm .           Tissue Specificity         Ubiquitous.           Function         function:Death-promoting transcriptional repressor "sequence caution:Contaminating sequence. Potential poly-A sequence, subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA. tissue specificity:Ubiquitous.           Background         This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing appotosis. Multiple	Specificity	BTF Monoclonal Antibody detects endogenous levels of BTF protein.
Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  BCLAF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf  Observed Band  106kD  Cell Pathway  Cytoplasm. Nucleus. Nucleus speckle. Nucleus, nucleoplasm.  Tissue Specificity  Ubiquitous.  Function  function:Death-promoting transcriptional repressor.,sequence caution:Contaminating sequence. Potential poly-A sequence.,subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA, tissue specificity: Ubiquitous.  Background  This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms BCLAF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf  Observed Band 106kD  Cell Pathway Cytoplasm. Nucleus. Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity Ubiquitous.  Function function:Death-promoting transcriptional repressor.,sequence caution:Contaminating sequence. Potential poly-A sequence.,subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA.,tissue specificity:Ubiquitous.,  Background This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of this protein induces apoptosis, which can be suppressed by co-expression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Source	Monoclonal, Mouse,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       BCLAF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf         Observed Band       106kD         Cell Pathway       Cytoplasm. Nucleus. Nucleus speckle . Nucleus, nucleoplasm .         Tissue Specificity       Ubiquitous.         Function       function:Death-promoting transcriptional repressor.,sequence caution:Contaminating sequence. Potential poly-A sequence.,subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA, tissue specificity:Ubiquitous.         Background       This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms BCLAF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf  Observed Band 106kD  Cell Pathway Cytoplasm. Nucleus. Nucleus speckle . Nucleus, nucleoplasm .  Tissue Specificity Ubiquitous.  Function function:Death-promoting transcriptional repressor.,sequence caution:Contaminating sequence. Potential poly-A sequence.,subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA.,tissue specificity:Ubiquitous.,  Background This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Dilution	WB 1:500-2000
Storage Stability  -20°C/1 year  Synonyms  BCLAF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf  Observed Band  106kD  Cell Pathway  Cytoplasm. Nucleus. Nucleus speckle. Nucleus, nucleoplasm.  Tissue Specificity  Ubiquitous.  Function  function:Death-promoting transcriptional repressor.,sequence caution:Contaminating sequence. Potential poly-A sequence.,subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA.,tissue specificity:Ubiquitous.,  This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Concentration	1 mg/ml
Synonyms  BCLAF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf  Observed Band  106kD  Cell Pathway  Cytoplasm. Nucleus. Nucleus speckle. Nucleus, nucleoplasm.  Tissue Specificity  Ubiquitous.  Function  function:Death-promoting transcriptional repressor.,sequence caution:Contaminating sequence. Potential poly-A sequence.,subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA.,tissue specificity:Ubiquitous.,  Background  This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Purity	≥90%
Observed Band  Cell Pathway  Cytoplasm. Nucleus. Nucleus speckle . Nucleus, nucleoplasm .  Ubiquitous.  Function  function: Death-promoting transcriptional repressor., sequence caution: Contaminating sequence. Potential poly-A sequence., subunit: Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA., tissue specificity: Ubiquitous.,  This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of this protein induces apoptosis, which can be suppressed by co-expression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Storage Stability	-20°C/1 year
Cell Pathway  Cytoplasm. Nucleus. Nucleus speckle . Nucleus, nucleoplasm .  Ubiquitous.  Function  function:Death-promoting transcriptional repressor.,sequence caution:Contaminating sequence. Potential poly-A sequence.,subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA.,tissue specificity:Ubiquitous.,  Background  This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of this protein induces apoptosis, which can be suppressed by co-expression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Synonyms	BCLAF1; BTF; KIAA0164; Bcl-2-associated transcription factor 1; Btf
Tissue Specificity  Function  function:Death-promoting transcriptional repressor.,sequence caution:Contaminating sequence. Potential poly-A sequence.,subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA.,tissue specificity:Ubiquitous.,  Background  This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of this protein induces apoptosis, which can be suppressed by co-expression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Observed Band	106kD
Function  function:Death-promoting transcriptional repressor.,sequence caution:Contaminating sequence. Potential poly-A sequence.,subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA.,tissue specificity:Ubiquitous.,  This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of this protein induces apoptosis, which can be suppressed by co-expression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Cell Pathway	Cytoplasm. Nucleus. Nucleus speckle . Nucleus, nucleoplasm .
caution:Contaminating sequence. Potential poly-A sequence., subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and with DNA., tissue specificity:Ubiquitous.,  Background  This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of this protein induces apoptosis, which can be suppressed by co-expression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Tissue Specificity	Ubiquitous.
members of the BCL2 family of proteins. Overexpression of this protein induces apoptosis, which can be suppressed by co-expression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple	Function	caution:Contaminating sequence. Potential poly-A sequence.,subunit:Interacts with Bcl-2 related proteins, EMD, with the adenovirus E1B 19 kDa protein and
	Background	members of the BCL2 family of proteins. Overexpression of this protein induces apoptosis, which can be suppressed by co-expression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple

Nanjing BYabscience technology Co.,Ltd





[provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using BTF Monoclonal Antibody