



## NY-CO-1 Monoclonal Antibody

Catalog No	BYmab-00466
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	NEMF
Protein Name	Nuclear export mediator factor NEMF
Immunogen	The antiserum was produced against synthesized peptide derived from human SDCG1. AA range:881-930
Specificity	NY-CO-1 Monoclonal Antibody detects endogenous levels of NY-CO-1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NEMF; SDCCAG1; Nuclear export mediator factor NEMF; Antigen NY-CO-1; Serologically defined colon cancer antigen 1
Observed Band	123kD
Cell Pathway	Cytoplasm, cytosol . Nucleus .
Tissue Specificity	Expressed in brain, heart, liver, lung, spleen, and skeletal muscle. Also expressed at lower levels in stomach and testis.
Function	
Background	This gene encodes a component of the ribosome quality control complex. The encoded protein facilitates the recognition and ubiquitination of stalled 60S subunits by the ubiquitin ligase listerin. A similar protein in fly functions as a tumor suppressor. [provided by RefSeq, Jul 2016],
matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



**Usage suggestions** 

(kD) 3T3

85-

48-

34-

26-

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

