



## PA26 Monoclonal Antibody

Catalog No	BYmab-00478
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SESN1
Protein Name	Sestrin-1
Immunogen	The antiserum was produced against synthesized peptide derived from human SESN1. AA range:271-320
Specificity	PA26 Monoclonal Antibody detects endogenous levels of PA26 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	SESN1; PA26; SEST1; Sestrin-1; p53-regulated protein PA26
Observed Band	57kD
Cell Pathway	Nucleus . Cytoplasm .
Tissue Specificity	Widely expressed.
Function	disease:Defects in SESN1 may be involved in heterotaxia. Heterotaxia is an aetiologically heterogeneous condition caused by an abnormal left-right axis formation, resulting in reversed left-right polarity of one or more organ systems.,function:Involved in the reduction of peroxiredoxins. May also be regulator of cellular growth.,induction:Isoforms T2 and isoform T3 are induced by genotoxic stress (UV, gamma-irradiation and cytotoxic drugs) in a p53-dependent manner. Isoform T1 is not induced by p53.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the sestrin family.,tissue specificity:Widely expressed.,

Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background	This gene encodes a member of the sestrin family. Sestrins are induced by the p53 tumor suppressor protein and play a role in the cellular response to DNA damage and oxidative stress. The encoded protein mediates p53 inhibition of cell growth by activating AMP-activated protein kinase, which results in the inhibition of the mammalian target of rapamycin protein. The encoded protein also plays a critical role in antioxidant defense by regenerating overoxidized peroxiredoxins, and the expression of this gene is a potential marker for exposure to radiation. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Dec 2010],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Nanjing BYabscience technology Co.,Ltd