



NOXIN Monoclonal Antibody

Catalog No	BYmab-00581
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	NOXIN
Protein Name	Nitric oxide-inducible gene protein
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human NOXIN. AA range:111-160
Specificity	NOXIN Monoclonal Antibody detects endogenous levels of NOXIN protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NOXIN; C11orf82; Nitric oxide-inducible gene protein
Observed Band	85kD
Cell Pathway	Cytoplasm . Nucleus . Accumulates in the nucleus in response to stress. .
Tissue Specificity	Highly expressed in colorectal and lung cancer tissues.
Function	function:Acts as an anti-apoptotic factor and its absence increases cell death under normal and stress conditions. Can induce cell cycle arrest in the G1 or early S phase and this activity is independent of TP53/p53.,subcellular location:Accumulates in the nucleus in response to stress.,
Background	function:Acts as an anti-apoptotic factor and its absence increases cell death under normal and stress conditions. Can induce cell cycle arrest in the G1 or early S phase and this activity is independent of TP53/p53.,subcellular location:Accumulates in the nucleus in response to stress.,

Nanjing BYabscience technology Co.,Ltd




matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using NOXIN Monoclonal Antibody