



## AI-BP Monoclonal Antibody

Catalog No	BYmab-00663
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	APOA1BP
Protein Name	NAD(P)H-hydrate epimerase
Immunogen	The antiserum was produced against synthesized peptide derived from human AIBP. AA range:121-170
Specificity	AI-BP Monoclonal Antibody detects endogenous levels of AI-BP protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
-	
Storage Stability	-20°C/1 year  APOA1BP; AIBP; YJEFN1; NAD(P)H-hydrate epimerase; Apolipoprotein A-I-binding protein; AI-BP; NAD(P)HX epimerase; YjeF N-terminal
Storage Stability Synonyms	-20°C/1 year  APOA1BP; AIBP; YJEFN1; NAD(P)H-hydrate epimerase; Apolipoprotein A-I-binding protein; AI-BP; NAD(P)HX epimerase; YjeF N-terminal domain-containing protein 1; YjeF_N1
Storage Stability Synonyms Observed Band	-20°C/1 year  APOA1BP; AIBP; YJEFN1; NAD(P)H-hydrate epimerase; Apolipoprotein A-I-binding protein; AI-BP; NAD(P)HX epimerase; YjeF N-terminal domain-containing protein 1; YjeF_N1 30kD  Mitochondrion . Secreted . In sperm, secretion gradually increases during
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year  APOA1BP; AIBP; YJEFN1; NAD(P)H-hydrate epimerase; Apolipoprotein A-I-binding protein; AI-BP; NAD(P)HX epimerase; YjeF N-terminal domain-containing protein 1; YjeF_N1  30kD  Mitochondrion . Secreted . In sperm, secretion gradually increases during capacitation  Ubiquitously expressed, with highest levels in kidney, heart and liver. Present in cerebrospinal fluid and urine but not in serum from healthy patients. Present in
Storage Stability Synonyms  Observed Band Cell Pathway Tissue Specificity	-20°C/1 year  APOA1BP; AIBP; YJEFN1; NAD(P)H-hydrate epimerase; Apolipoprotein A-I-binding protein; AI-BP; NAD(P)HX epimerase; YjeF N-terminal domain-containing protein 1; YjeF_N1 30kD  Mitochondrion . Secreted . In sperm, secretion gradually increases during capacitation  Ubiquitously expressed, with highest levels in kidney, heart and liver. Present in cerebrospinal fluid and urine but not in serum from healthy patients. Present in serum of sepsis patients (at protein level).  similarity:Contains 1 YjeF N-terminal domain.,subunit:Interacts with APOA1 and APOA2.,tissue specificity:Ubiquitously expressed, with highest levels in kidney, heart and liver. Present in cerebrospinal fluid and urine but not in serum from


Nanjing BYabscience technology Co.,Ltd





	with apoA-I. The human genome contains related pseudogenes. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using Al-BP Monoclonal Antibody

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658