



GS28 Monoclonal Antibody

Catalog No	BYmab-00703
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	GOSR1
Protein Name	Golgi SNAP receptor complex member 1
Immunogen	The antiserum was produced against synthesized peptide derived from human GOSR1. AA range:11-60
Specificity	GS28 Monoclonal Antibody detects endogenous levels of GS28 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	GOSR1; GS28; Golgi SNAP receptor complex member 1; 28 kDa Golgi SNARE protein; 28 kDa cis-Golgi SNARE p28; GOS-28
Synonyms Observed Band	·
	protein; 28 kDa cis-Golgi SNARE p28; GOS-28
Observed Band	protein; 28 kDa cis-Golgi SNARE p28; GOS-28 28kD Golgi apparatus membrane; Single-pass type IV membrane protein. Localizes throughout the Golgi apparatus, with lowest levels in the trans-Golgi network (By similarity). Enriched on vesicular components at the terminal rims of the Golgi.
Observed Band Cell Pathway	protein; 28 kDa cis-Golgi SNARE p28; GOS-28 28kD Golgi apparatus membrane; Single-pass type IV membrane protein. Localizes throughout the Golgi apparatus, with lowest levels in the trans-Golgi network (By similarity). Enriched on vesicular components at the terminal rims of the Golgi. Found in Golgi microtubules at low temperature (15 degrees Celsius). Colon carcinoma,Prostate,Testis,Umbilical cord blood, function:Involved in transport from the ER to the Golgi apparatus as well as in intra-Golgi transport.,similarity:Belongs to the GOSR1 family.,subcellular location:Enriched on vesicular components at the terminal rims of the
Observed Band Cell Pathway Tissue Specificity	protein; 28 kDa cis-Golgi SNARE p28; GOS-28 28kD Golgi apparatus membrane; Single-pass type IV membrane protein. Localizes throughout the Golgi apparatus, with lowest levels in the trans-Golgi network (By similarity). Enriched on vesicular components at the terminal rims of the Golgi. Found in Golgi microtubules at low temperature (15 degrees Celsius). Colon carcinoma,Prostate,Testis,Umbilical cord blood, function:Involved in transport from the ER to the Golgi apparatus as well as in intra-Golgi transport.,similarity:Belongs to the GOSR1 family.,subcellular location:Enriched on vesicular components at the terminal rims of the Golgi.,subunit:Interacts with GABARAPL2. Identified in a unique SNARE complex

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





	compartments. This protein is considered an essential component of the Golgi SNAP receptor (SNARE) complex. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

