



NRBF-2 Monoclonal Antibody

Catalog No BYmab-00706 Isotype IgG Reactivity Human;Mouse;Rat Applications WB Gene Name NRBF2 Protein Name Nuclear receptor-binding factor 2 Immunogen The antiserum was produced against synthesized peptide derived from human NRBF2. An range;140-189 Specificity NRBF-2 Monoclonal Antibody detects endogenous levels of NRBF-2 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome subset of cells in pediatric medulioblastoma (PubMed:15610520). Expressed in a subset of cells in pediatric medulioblastoma (PubMed:18619852). Function function:May modulate transcri		
Reactivity Human;Mouse;Rat Applications WB Gene Name NRBF2 Protein Name Nuclear receptor-binding factor 2 Immunogen The antiserum was produced against synthesized peptide derived from human NRBF2. AA range:140-189 Specificity NRBF-2 Monoclonal Antibody detects endogenous levels of NRBF-2 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR Observed Band 36kD Cell Pathway Nucleus . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome . Tissue Specificity Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound	Catalog No	BYmab-00706
Applications Gene Name NRBF2 Protein Name Nuclear receptor-binding factor 2 Immunogen The antiserum was produced against synthesized peptide derived from human NRBF2. AA range:140-189 Specificity NRBF-2 Monoclonal Antibody detects endogenous levels of NRBF-2 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR Observed Band 36kD Cell Pathway Nucleus. Cytoplasm. Cytoplasmic vesicle. Cytoplasmic vesicle, autophagosome Tissue Specificity Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activator in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand, tissue specificity:Detected in keratinocytes, liver and placenta. Function:May modulate transcriptional activator by target nuclear receptors. Can act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound	Isotype	IgG
Protein Name NRBF2 Protein Name Nuclear receptor-binding factor 2 Immunogen The antiserum was produced against synthesized peptide derived from human NRBF2. AA range:140-189 Specificity NRBF-2 Monoclonal Antibody detects endogenous levels of NRBF-2 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR Observed Band 36kD Cell Pathway Nucleus . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome Tissue Specificity Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activator in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand, tissue specificity:Detected in keratinocytes, liver and placenta. Background function:May modulate transcriptional activator by target nuclear receptors. Can act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG	Reactivity	Human;Mouse;Rat
Protein Name Nuclear receptor-binding factor 2 Immunogen The antiserum was produced against synthesized peptide derived from human NRBF2. AA range:140-189 Specificity NRBF-2 Monoclonal Antibody detects endogenous levels of NRBF-2 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome . Subset of cells in pediatric medulloblastoma (PubMed:18619852). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Expressed in and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand, lissue specificity: Detected in keratinocytes, liver and placenta. Background function: May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro). subunit.Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bou	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human NRBF2. AA range:140-189 Specificity NRBF-2 Monoclonal Antibody detects endogenous levels of NRBF-2 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome Tissue Specificity Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound largend, tissue specificity:Detected in keratinocytes, lisuer, subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RA	Gene Name	NRBF2
NRBF2. AA range:140-189 Specificity NRBF-2 Monoclonal Antibody detects endogenous levels of NRBF-2 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR Observed Band 36kD Cell Pathway Nucleus . Cytoplasmic Vesicle . Cytoplasmic vesicle, autophagosome Tissue Specificity Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand, tissue specificity:Detected in keratinocytes, liver and placenta., Background function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with PPARA, and the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with PPARA, and the presence of bound and PPARG. Interacts with PPARA, and the presence of bound and PPARG. Interacts with PPARA, and the presence of bound and PPARG. Interacts with PPARA, and the presence of bound and PPARG. Interacts with PPARA, and the presence of bound and PPARG. Interacts with PPARA, and the presence of bound and PPARG. Interacts with PPARA and the PPARB.	Protein Name	Nuclear receptor-binding factor 2
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome Tissue Specificity Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand, tissue specificity:Detected in keratinocytes, liver and placenta. Background function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and PRARG interpresence of bound	lmmunogen	
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR Observed Band 36kD Cell Pathway Nucleus . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome Tissue Specificity Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG.	Specificity	NRBF-2 Monoclonal Antibody detects endogenous levels of NRBF-2 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR Observed Band 36kD Cell Pathway Nucleus . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome . Tissue Specificity Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome . Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand., tissue specificity:Detected in keratinocytes, liver and placenta. Background function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome . Tissue Specificity Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand.,tissue specificity:Detected in keratinocytes, liver and placenta., Background function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome . Tissue Specificity Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand.,tissue specificity:Detected in keratinocytes, liver and placenta., Background function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome . Tissue Specificity Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand.,tissue specificity:Detected in keratinocytes, liver and placenta., Background function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound	Concentration	1 mg/ml
Synonyms NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome . Tissue Specificity Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand.,tissue specificity:Detected in keratinocytes, liver and placenta., function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound	Purity	≥90%
PPAR and RXR 36kD Cell Pathway Nucleus . Cytoplasm . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome . Tissue Specificity Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand.,tissue specificity:Detected in keratinocytes, liver and placenta., function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound	Storage Stability	-20°C/1 year
Cell Pathway Nucleus . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand.,tissue specificity:Detected in keratinocytes, liver and placenta., function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound	Synonyms	. •
Tissue Specificity Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand.,tissue specificity:Detected in keratinocytes, liver and placenta., function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound	Observed Band	36kD
subset of cells in pediatric medulloblastoma (PubMed:18619852). Function function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand.,tissue specificity:Detected in keratinocytes, liver and placenta., function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound	Cell Pathway	Nucleus . Cytoplasm . Cytoplasmic vesicle . Cytoplasmic vesicle, autophagosome .
act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound ligand.,tissue specificity:Detected in keratinocytes, liver and placenta., Background function:May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound	Tissue Specificity	Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852).
act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARA, and PPARA, Interacts with RARA, RARG and RXRA in the presence of bound	Function	act as transcriptional activator (in vitro), subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound
	Background	act as transcriptional activator (in vitro).,subunit:Interacts with PPARA, PPARD and PPARG. Interacts with RARA, RARG and RXRA in the presence of bound

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658






matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using NRBF-2 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd