



# ApoC-III Monoclonal Antibody

<b>Catalog No</b>	BYmab-00774
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	APOC3
<b>Protein Name</b>	Apolipoprotein C-III
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from the C-terminal region of human APOC3. AA range:46-95
<b>Specificity</b>	ApoC-III Monoclonal Antibody detects endogenous levels of ApoC-III protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	APOC3; Apolipoprotein C-III; Apo-CIII; ApoC-III; Apolipoprotein C3
<b>Observed Band</b>	11kD
<b>Cell Pathway</b>	Secreted .
<b>Tissue Specificity</b>	Liver.
<b>Function</b>	disease:Defects in APOC3 may be a cause of hyperalphalipoproteinemia [MIM:143470]. Affected individuals show high levels of alpha-lipoprotein (high density lipoprotein/HDL).,function:Inhibits lipoprotein lipase and hepatic lipase and decreases the uptake of lymph chylomicrons by hepatic cells. This suggests that it delays the catabolism of triglyceride-rich particles.,PTM:O-linked glycan consists of Gal-GalNAc disaccharide, further modified with up to 3 sialic acid residues.,similarity:Belongs to the apolipoprotein C3 family.,tissue specificity:Constitutes 50% of the protein fraction of VLDL and 2% of that of HDL. Synthesized predominantly in liver and to a lesser degree in intestine.,

**Nanjing BYabscience technology Co.,Ltd**

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658



## Background

Apolipoprotein C-III is a very low density lipoprotein (VLDL) protein. APOC3 inhibits lipoprotein lipase and hepatic lipase; it is thought to delay catabolism of triglyceride-rich particles. The APOA1, APOC3 and APOA4 genes are closely linked in both rat and human genomes. The A-I and A-IV genes are transcribed from the same strand, while the A-I and C-III genes are convergently transcribed. An increase in apoC-III levels induces the development of hypertriglyceridemia. [provided by RefSeq, Jul 2008],


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using ApoC-III Monoclonal Antibody