



Elk-1 (phospho Ser383) Monoclonal Antibody

Catalog No	BYmab-01237
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ELK1
Protein Name	ETS domain-containing protein Elk-1
Immunogen	The antiserum was produced against synthesized peptide derived from human Elk1 around the phosphorylation site of Ser383. AA range:351-400
Specificity	Phospho-Elk-1 (S383) Monoclonal Antibody detects endogenous levels of Elk-1 protein only when phosphorylated at S383.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ELK1; ETS domain-containing protein Elk-1
Observed Band	
Cell Pathway	Nucleus.
Tissue Specificity	Lung and testis.
Function	function:Stimulates transcription. Binds to purine-rich DNA sequences. Can form a ternary complex with the serum response factor and the ETS and SRF motifs of the fos serum response element.,PTM:On mitogenic stimulation, phosphorylated on C-terminal serine and threonine residues by MAPK1. Ser-383 and Ser-389 are the preferred sites for MAPK1. In vitro, phosphorylation by MAPK1 potentiates ternary complex formation with the serum responses factors, SRE and SRF. Phosphorylation leads to loss of sumoylation and restores transcriptional activator activity.,PTM:Sumoylation represses transcriptional activator activity as it results in recruitment of HDAC2 to target gene promoters which leads to decreased histone acetylation and reduced transactivator activity. It also regulates nuclear retention.,similarity:Belongs to the ETS family.,similarity:Contains 1 ETS

Nanjing BYabscience technology Co.,Ltd



DNA-binding domain.,subunit:Intera

Background

This gene is a member of the Ets family of transcription factors and of the ternary complex factor (TCF) subfamily. Proteins of the TCF subfamily form a ternary complex by binding to the the serum response factor and the serum response element in the promoter of the c-fos proto-oncogene. The protein encoded by this gene is a nuclear target for the ras-raf-MAPK signaling cascade. This gene produces multiple isoforms by using alternative translational start codons and by alternative splicing. Related pseudogenes have been identified on chromosomes 7 and 14. [provided by RefSeq, Mar 2012],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Elk-1 (phospho Ser383) Monoclonal Antibody

Nanjing BYabs science technology Co.,Ltd