



Jun D (phospho Ser255) Monoclonal Antibody

Catalog No BYmab-01258 Isotype IgG Reactivity Human:Mouse;Rat Applications WB Gene Name JUND Protein Name Transcription factor jun-D Immunogen The antiserum was produced against synthesized peptide derived from human JunD around the phosphorylation site of Ser255. An range:222-271 Specificity Phospho-Jun D (S255) Monoclonal Antibody detects endogenous levels of Jun D protein only when phosphorylated at S255. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms JUND; Transcription factor jun-D Observed Band Cell Pathway Nucleus. Tissue Specificity Epithelium, Spleen, Function function:Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site, similarity, Belongs to the bZIP familly. Julusulfa		
Reactivity Human;Mouse;Rat Applications WB Gene Name JUND Protein Name Transcription factor jun-D Immunogen The antiserum was produced against synthesized peptide derived from human JunD around the phosphorylation site of Ser255. AA range:222-271 Specificity Phospho-Jun D (S255) Monoclonal Antibody detects endogenous levels of Jun D protein only when phosphorylated at \$255. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms JUND; Transcription factor jun-D Observed Band Cell Pathway Nucleus. Tissue Specificity Epithelium, Spleen, Function function:Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site, similarity:Belongs to the bZIP family, Jun subfamily, similarity:Contains 1 bZIP domain, subunit:Binds DNA as a dimer., Background The pr	Catalog No	BYmab-01258
Applications WB Gene Name JUND Protein Name Transcription factor jun-D Immunogen The antiserum was produced against synthesized peptide derived from human JunD around the phosphorylation site of Ser255. AA range:222-271 Specificity Phospho-Jun D (S255) Monoclonal Antibody detects endogenous levels of Jun D protein only when phosphorylated at S255. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms JUND; Transcription factor jun-D Observed Band Cell Pathway Nucleus. Tissue Specificity Epithelium,Spleen, Function function:Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site, similarity; Belongs to the bZIP family. Jun subfamily, similarity:Contains 1 bZIP domain., subunit:Binds DNA as a dimer., The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Isotype	IgG
Gene Name Protein Name Transcription factor jun-D Immunogen The antiserum was produced against synthesized peptide derived from human JunD around the phosphorylation site of Ser255. AA range:222-271 Specificity Phospho-Jun D (S255) Monoclonal Antibody detects endogenous levels of Jun D protein only when phosphorylated at S255. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms JUND; Transcription factor jun-D Observed Band Cell Pathway Nucleus. Tissue Specificity Epithelium,Spleen, function:Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site, similarity:Belongs to the bZIP family. Jun subfamily, similarity:Contains 1 bZIP domain, subunit:Binds DNA as a dimer, a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Reactivity	Human;Mouse;Rat
Protein Name Transcription factor jun-D Immunogen The antiserum was produced against synthesized peptide derived from human JunD around the phosphorylation site of Ser255. AA range:222-271 Specificity Phospho-Jun D (S255) Monoclonal Antibody detects endogenous levels of Jun D protein only When phosphorylated at S255. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms JUND; Transcription factor jun-D Observed Band Cell Pathway Cell Pathway Nucleus. Tissue Specificity Epithelium,Spleen, Function function:Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site, similarity:Belongs to the bZIP family. Jun subfamily, similarity:Contains 1 bZIP domain, subunit:Binds DNA as a dimer, subfamily, similarity:Contains 1 bZIP domain, subunit:Binds DNA as a dimer, a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alterna	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human JunD around the phosphorylation site of Ser255. AA range:222-271 Specificity Phospho-Jun D (S255) Monoclonal Antibody detects endogenous levels of Jun D protein only when phosphorylated at S255. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms JUND; Transcription factor jun-D Observed Band Cell Pathway Nucleus. Tissue Specificity Epithelium, Spleen, function: Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site, similarity; Belongs to the bZIP family. Jun subfamily, similarity. Contains 1 bZIP domain, subunit: Binds DNA as a dimer., Background The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Gene Name	JUND
JunD around the phosphorylation site of Ser255. AA range:222-271 Specificity Phospho-Jun D (S255) Monoclonal Antibody detects endogenous levels of Jun D protein only when phosphorylated at S255. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms JUND; Transcription factor jun-D Observed Band Cell Pathway Nucleus. Tissue Specificity Epithelium,Spleen, Function function:Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site, similarity:Belongs to the bZIP family. Jun subfamily., similarity:Contains 1 bZIP domain., subunit:Binds DNA as a dimer., Background The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Protein Name	Transcription factor jun-D
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms JUND; Transcription factor jun-D Observed Band Cell Pathway Nucleus. Tissue Specificity Epithelium, Spleen, Function function: Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site., similarity: Belongs to the bZIP family. Jun subfamily., similarity: Contains 1 bZIP domain., subunit: Binds DNA as a dimer., Background The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Immunogen	
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms JUND; Transcription factor jun-D Observed Band Cell Pathway Nucleus. Tissue Specificity Epithelium, Spleen, Function function: Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site., similarity: Belongs to the bZIP family. Jun subfamily., similarity: Contains 1 bZIP domain., subunit: Binds DNA as a dimer., Background The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Specificity	Phospho-Jun D (S255) Monoclonal Antibody detects endogenous levels of Jun D protein only when phosphorylated at S255.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms JUND; Transcription factor jun-D Observed Band Cell Pathway Nucleus. Tissue Specificity Epithelium,Spleen, Function function:Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site.,similarity:Belongs to the bZIP family. Jun subfamily.,similarity:Contains 1 bZIP domain.,subunit:Binds DNA as a dimer., Background The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Formulation	
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms JUND; Transcription factor jun-D Observed Band Cell Pathway Nucleus. Tissue Specificity Epithelium,Spleen, Function function:Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site, similarity:Belongs to the bZIP family. Jun subfamily.,similarity:Contains 1 bZIP domain.,subunit:Binds DNA as a dimer., Background The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms JUND; Transcription factor jun-D Observed Band Cell Pathway Nucleus. Fissue Specificity Epithelium,Spleen, Function function:Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site, similarity:Belongs to the bZIP family. Jun subfamily., similarity:Contains 1 bZIP domain., subunit:Binds DNA as a dimer., Background The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Purification	•
Purity ≥90% Storage Stability -20°C/1 year Synonyms JUND; Transcription factor jun-D Observed Band Cell Pathway Nucleus. Tissue Specificity Epithelium, Spleen, Function function: Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site., similarity: Belongs to the bZIP family. Jun subfamily., similarity: Contains 1 bZIP domain., subunit: Binds DNA as a dimer., Background The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms JUND; Transcription factor jun-D Observed Band Cell Pathway Nucleus. Tissue Specificity Epithelium, Spleen, Function function: Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site., similarity: Belongs to the bZIP family. Jun subfamily., similarity: Contains 1 bZIP domain., subunit: Binds DNA as a dimer., Background The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Concentration	1 mg/ml
Synonyms JUND; Transcription factor jun-D Observed Band Cell Pathway Nucleus. Tissue Specificity Epithelium, Spleen, function: Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site., similarity: Belongs to the bZIP family. Jun subfamily., similarity: Contains 1 bZIP domain., subunit: Binds DNA as a dimer., Background The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Purity	≥90%
Observed Band Cell Pathway Nucleus. Tissue Specificity Epithelium, Spleen, function: Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site., similarity: Belongs to the bZIP family. Jun subfamily., similarity: Contains 1 bZIP domain., subunit: Binds DNA as a dimer., Background The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Storage Stability	-20°C/1 year
Cell Pathway Nucleus. Tissue Specificity Epithelium, Spleen, Function function:Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site., similarity:Belongs to the bZIP family. Jun subfamily., similarity:Contains 1 bZIP domain., subunit:Binds DNA as a dimer., Background The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Synonyms	JUND; Transcription factor jun-D
Tissue Specificity Epithelium, Spleen, function function: Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site., similarity: Belongs to the bZIP family. Jun subfamily., similarity: Contains 1 bZIP domain., subunit: Binds DNA as a dimer., The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Observed Band	
Function function:Binds an AP-1 site and upon cotransfection stimulates the activity of a promoter that bears an AP-1 site.,similarity:Belongs to the bZIP family. Jun subfamily.,similarity:Contains 1 bZIP domain.,subunit:Binds DNA as a dimer., The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Cell Pathway	Nucleus.
promoter that bears an AP-1 site., similarity: Belongs to the bZIP family. Jun subfamily., similarity: Contains 1 bZIP domain., subunit: Binds DNA as a dimer., The protein encoded by this intronless gene is a member of the JUN family, and a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Tissue Specificity	Epithelium,Spleen,
a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different	Function	promoter that bears an AP-1 site. similarity:Belongs to the bZIP family. Jun
	Background	a functional component of the AP1 transcription factor complex. This protein has been proposed to protect cells from p53-dependent senescence and apoptosis. Alternative translation initiation site usage results in the production of different

Nanjing BYabscience technology Co.,Ltd





matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

