



hnRNP K (phospho Ser216) Monoclonal Antibody

Catalog No	BYmab-01328
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	HNRNPK
Protein Name	Heterogeneous nuclear ribonucleoprotein K
Immunogen	Synthesized phospho-peptide around the phosphorylation site of human hnRNP K (phospho Ser216)
Specificity	Phospho-hnRNP K (S216) Monoclonal Antibody detects endogenous levels of hnRNP K protein only when phosphorylated at S216.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HNRNPK; HNRPK; Heterogeneous nuclear ribonucleoprotein K; hnRNP K; Transformation up-regulated nuclear protein; TUNP
Observed Band	55kD
Cell Pathway	Cytoplasm . Nucleus, nucleoplasm . Cell projection, podosome . Recruited to p53/TP53-responsive promoters, in the presence of functional p53/TP53 (PubMed:16360036). In case of ASFV infection, there is a shift in the localization which becomes predominantly nuclear (PubMed:18775702).
Tissue Specificity	Brain,Cajal-Retzius cell,Colorectal cancer and surrounding
Function	function:One of the major pre-mRNA-binding proteins. Binds tenaciously to poly(C) sequences. Likely to play a role in the nuclear metabolism of hnRNAs, particularly for pre-mRNAs that contain cytidine-rich sequences. Can also bind poly(C) single-stranded DNA.,mass spectrometry: PubMed:11840567,PTM:Arg-296 and Arg-299 are dimethylated, probably to asymmetric dimethylarginine.,similarity:Contains 1 KH domain.,similarity:Contains 2 KH domains.,similarity:Contains 3 KH domains.,subcellular location:In case of ASFV infection, there is a shift in the localization which becomes predominantly

Nanjing BYabscience technology Co.,Ltd



nuclear.,subunit:Interacts with RBM42 and ZIK1 (By similarity). Identified in the spliceosome C complex, at least composed of AQR, ASCC3L1, C19orf29, CDC40, CDC5L, CRNKL1, DDX23, DDX41, DDX48, DDX5, DGCR14, DHX35, DHX38, DHX8, EFTUD2, FRG1, GPATC1, HNRPA1, HNRPA2B1, HNRPA3, HNRPC, HNRPF, HNRPH1, HNRNPK, HNR

Background

This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene is located in the nucleoplasm and has three repeats of KH domains that binds to RNAs. It is distinct among other hnRNP proteins in its binding preference; it binds tenaciously to poly(C). This protein is also thought to have a role during cell cycle progression. Several alternatively spliced transcript variants have

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images