



# NFκB-p105 (phospho Ser927) Monoclonal Antibody

<b>Catalog No</b>	BYmab-01383
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	NFKB1
<b>Protein Name</b>	Nuclear factor NF-kappa-B p105 subunit
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human NF-kappaB p105/p50 around the phosphorylation site of Ser927. AA range:896-945
<b>Specificity</b>	Phospho-NF κ B-p105 (S927) Monoclonal Antibody detects endogenous levels of NF κ B-p105 protein only when phosphorylated at S927.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	NFKB1; Nuclear factor NF-kappa-B p105 subunit; DNA-binding factor KBF1; EBP-1; Nuclear factor of kappa light polypeptide gene enhancer in B-cells 1
<b>Observed Band</b>	110kD
<b>Cell Pathway</b>	Nucleus. Cytoplasm. Nuclear, but also found in the cytoplasm in an inactive form complexed to an inhibitor (I-kappa-B).
<b>Tissue Specificity</b>	Muscle,Rectum tumor,Uterus,
<b>Function</b>	domain:Glycine-rich region (GRR) appears to be a critical element in the generation of p50.;domain:The C-terminus of p105 might be involved in cytoplasmic retention, inhibition of DNA-binding, and transcription activation.;function:NF-kappa-B is a pleiotropic transcription factor which is present in almost all cell types and is involved in many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50,

**Nanjing BYabscience technology Co.,Ltd**



REL and NF $\kappa$ B2/p52 and the heterodimeric p65-p50 complex appears to be most abundant one. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Diff

## Background

nuclear factor kappa B subunit 1(NF $\kappa$ B1) Homo sapiens This gene encodes a 105 kD protein which can undergo cotranslational processing by the 26S proteasome to produce a 50 kD protein. The 105 kD protein is a Rel protein-specific transcription inhibitor and the 50 kD protein is a DNA binding subunit of the NF-kappa-B (NF $\kappa$ B) protein complex. NF $\kappa$ B is a transcription regulator that is activated by various intra- and extra-cellular stimuli such as cytokines, oxidant-free radicals, ultraviolet irradiation, and bacterial or viral products. Activated NF $\kappa$ B translocates into the nucleus and stimulates the expression of genes involved in a wide variety of biological functions. Inappropriate activation of NF $\kappa$ B has been associated with a number of inflammatory diseases while persistent inhibition of NF $\kappa$ B leads to inappropriate immune cell development or delayed cell growth. Alternative splicing results in multiple transcript variants encoding different isof


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using NF  $\kappa$  B-p105 (phospho Ser927) Monoclonal Antibody

Nanjing BYabs science technology Co.,Ltd