



A20 Monoclonal Antibody

Catalog No	BYmab-01516
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	TNFAIP3
Protein Name	Tumor necrosis factor alpha-induced protein 3
Immunogen	The antiserum was produced against synthesized peptide derived from human TNAP3. AA range:321-370
Specificity	A20 Monoclonal Antibody detects endogenous levels of A20 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TNFAIP3; OTUD7C; Tumor necrosis factor alpha-induced protein 3; TNF alpha-induced protein 3; OTU domain-containing protein 7C; Putative DNA-binding protein A20; Zinc finger protein A20
Observed Band	89kD
Cell Pathway	Cytoplasm. Nucleus. Lysosome.; [A20p50]: Cytoplasm.
Tissue Specificity	Epithelium,Testis,Trachea,
Function	function:Interacts with NAF1 and inhibits TNF-induced NF-kappa-B-dependent gene expression by interfering with an RIP- or TRAF2-mediated transactivation signal. Inhibitor of programmed cell death. Has a role in the function of the lymphoid system and may contribute to the in vivo effects of TNF (By similarity). Has deubiquitinating activity that is directed towards 'Lys-48' or 'Lys-63'-linked polyubiquitin chains.,induction:By TNF-alpha.,similarity:Belongs to the peptidase C64 family.,similarity:Contains 1 OTU domain.,similarity:Contains 7 A20-type zinc fingers.,subunit:Homodimer. Interacts with NAF1, TAX1BP1 and TRAF2.,

Nanjing BYabscience technology Co.,Ltd



Background

This gene was identified as a gene whose expression is rapidly induced by the tumor necrosis factor (TNF). The protein encoded by this gene is a zinc finger protein and ubiquitin-editing enzyme, and has been shown to inhibit NF-kappa B activation as well as TNF-mediated apoptosis. The encoded protein, which has both ubiquitin ligase and deubiquitinase activities, is involved in the cytokine-mediated immune and inflammatory responses. Several transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2012],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using A20 Monoclonal Antibody