



ASC-1 Monoclonal Antibody

Catalog No	BYmab-01540
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	TRIP4
Protein Name	Activating signal cointegrator 1
Immunogen	The antiserum was produced against synthesized peptide derived from human TRIP4. AA range:81-130
Specificity	ASC-1 Monoclonal Antibody detects endogenous levels of ASC-1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TRIP4; Activating signal cointegrator 1; ASC-1; Thyroid receptor-interacting protein 4; TR-interacting protein 4; TRIP-4
Observed Band	66kD
Cell Pathway	Nucleus . Cytoplasm, cytosol . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasmic under conditions of serum deprivation (PubMed:10454579). Colocalizes with NEK6 in the centrosome (PubMed:20873783). .
Tissue Specificity	Lung,
Function	function:Transcription coactivator of nuclear receptors which functions in conjunction with CBP-p300 and SRC-1 and may play an important role in establishing distinct coactivator complexes under different cellular conditions. Plays a pivotal role in the transactivation of NF-kappa-B, SRF and AP1. Acts as a mediator of transrepression between nuclear receptor and either AP1 or NF-kappa-B. Plays a role in androgen receptor transactivation and in testicular function.,subcellular location:Cytoplasmic under conditions of serum deprivation.,subunit:Specifically interacts with the ligand binding domain of the

Nanjing BYabscience technology Co.,Ltd



thyroid receptor (TR). This interaction requires the presence of thyroid hormone. Exists as a steady-state complex associated with ASCC1, ASCC2 and HELIC1. Interacts with the androgen receptor androgen (AR) in an androgen, testosterone and dihydrotestosterone-dependent manner.,

Background

This gene encodes a subunit of the tetrameric nuclear activating signal cointegrator 1 (ASC-1) complex, which associates with transcriptional coactivators, nuclear receptors and basal transcription factors to facilitate nuclear receptors-mediated transcription. This protein is localized in the nucleus and contains an E1A-type zinc finger domain, which mediates interaction with transcriptional coactivators and ligand-bound nuclear receptors, such as thyroid hormone receptor and retinoid X receptor alpha, but not glucocorticoid receptor. Mutations in this gene are associated with spinal muscular atrophy with congenital bone fractures-1 (SMABF1). [provided by RefSeq, Apr 2016],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabs science technology Co.,Ltd