



## **ASF1B Monoclonal Antibody**

	DV 1 04540
Catalog No	BYmab-01543
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ASF1B
Protein Name	Histone chaperone ASF1B
Immunogen	The antiserum was produced against synthesized peptide derived from human ASF1B. AA range:101-150
Specificity	ASF1B Monoclonal Antibody detects endogenous levels of ASF1B protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ASF1B; Histone chaperone ASF1B; Anti-silencing function protein 1 homolog B; hAsf1; hAsf1b; CCG1-interacting factor A-II; CIA-II; hCIA-II
Observed Band	26kD
Cell Pathway	Nucleus .
Tissue Specificity	Highly expressed in testis and at lower levels in colon, small intestine and thymus.
Function	function: Histone chaperone that facilitates histone deposition and histone exchange and removal during nucleosome assembly and disassembly. Cooperates with chromatin assembly factor 1 (CAF-1) to promote replication-dependent chromatin assembly. Does not participate in replication-independent nucleosome deposition which is mediated by ASF1A and HIRA. Required for spermatogenesis.,PTM:Phosphorylated by TLK1 and TLK2.,similarity:Belongs to the ASF1 family.,subunit:Interacts with histone H3 (including both histone H3.1 and H3.3) and histone H4. Interacts with the CHAF1A, CHAF1B and RBBP4 subunits of the CAF-1 complex. Interacts with HAT1, NASP, TAF1, TLK1 and TLK2.,tissue specificity:Highly expressed in testis

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



and at lower levels in colon	, small intestine and thymus.,
	, errian niteetine arra trijinaer,

Background	This gene encodes a member of the H3/H4 family of histone chaperone proteins and is similar to the anti-silencing function-1 gene in yeast. The encoded protein is the substrate of the tousled-like kinase family of cell cycle-regulated kinases, and may play a key role in modulating the nucleosome structure of chromatin by
	may play a key role in modulating the nucleosome structure of chromatin by
	ensuring a constant supply of histones at sites of nucleosome assembly.

[provided by RefSeq, Jul 2008],

matters needing attention

**Usage suggestions** 

Avoid repeated freezing and thawing!

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



