



CNOT4 Monoclonal Antibody


| | |
|--------------------|--|
| Catalog No | BYmab-01616 |
| Isotype | IgG |
| Reactivity | Human;Mouse;Rat |
| Applications | WB |
| Gene Name | CNOT4 |
| Protein Name | CCR4-NOT transcription complex subunit 4 |
| Immunogen | The antiserum was produced against synthesized peptide derived from human CNOT4. AA range:61-110 |
| Specificity | CNOT4 Monoclonal Antibody detects endogenous levels of CNOT4 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | CNOT4; NOT4; CCR4-NOT transcription complex subunit 4; CCR4-associated factor 4; E3 ubiquitin-protein ligase CNOT4; Potential transcriptional repressor NOT4Hp |
| Observed Band | 64kD |
| Cell Pathway | Cytoplasm . Nucleus . |
| Tissue Specificity | Mammary gland,Testis, |
| Function | function:Has E3 ubiquitin ligase activity. The CCR4-NOT complex functions as general transcription regulation complex.,pathway:Protein modification; protein ubiquitination.,PTM:Autoubiquitinated.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,sequence caution:Sequencing errors.,similarity:Contains 1 C3H1-type zinc finger.,similarity:Contains 1 RING-type zinc finger.,similarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Subunit of the CCR4-NOT core complex that contains CHAF1A, CHAF1B, CNOT1, CNOT2, CNOT3, CNOT4, CNOT6 and CNOT8. Binds CNOT1 via its C-terminus. Binds E2 ubiquitin ligases via its RING domain., |

Nanjing BYabscience technology Co.,Ltd



| | |
|----------------------------------|--|
| Background | CCR4-NOT transcription complex subunit 4(CNOT4) Homo sapiens The protein encoded by this gene is a subunit of the CCR4-NOT complex, a global transcriptional regulator. The encoded protein interacts with CNOT1 and has E3 ubiquitin ligase activity. Several transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jul 2010], |
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |

Products Images



Nanjing BYabs science technology Co.,Ltd