



EF-1 γ Monoclonal Antibody

Catalog No	BYmab-01680
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	EEF1G
Protein Name	Elongation factor 1-gamma
Immunogen	The antiserum was produced against synthesized peptide derived from human EEF1G. AA range:101-150
Specificity	EF-1 γ Monoclonal Antibody detects endogenous levels of EF-1 γ protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Purity Storage Stability Synonyms	≥90% -20°C/1 year EEF1G; EF1G; Elongation factor 1-gamma; EF-1-gamma; eEF-1B gamma
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year EEF1G; EF1G; Elongation factor 1-gamma; EF-1-gamma; eEF-1B gamma 55kD nucleus,cytoplasm,endoplasmic reticulum,cytosol,cell-cell adherens
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year EEF1G; EF1G; Elongation factor 1-gamma; EF-1-gamma; eEF-1B gamma 55kD nucleus,cytoplasm,endoplasmic reticulum,cytosol,cell-cell adherens junction,membrane,extracellular exosome, Highly expressed in pancreatic tumor tissue and to a lesser extent in normal kidney, intestine, pancreas, stomach, lung, brain, spleen and liver. function:Probably plays a role in anchoring the complex to other cellular components.,similarity:Contains 1 EF-1-gamma C-terminal domain.,similarity:Contains 1 GST C-terminal domain.,similarity:Contains 1 GST N-terminal domain.,subunit:EF-1 is composed of four subunits: alpha, beta, delta, and gamma.,tissue specificity:Highly expressed in pancreatic tumor tissue and to
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year EEF1G; EF1G; Elongation factor 1-gamma; EF-1-gamma; eEF-1B gamma 55kD nucleus,cytoplasm,endoplasmic reticulum,cytosol,cell-cell adherens junction,membrane,extracellular exosome, Highly expressed in pancreatic tumor tissue and to a lesser extent in normal kidney, intestine, pancreas, stomach, lung, brain, spleen and liver. function:Probably plays a role in anchoring the complex to other cellular components.,similarity:Contains 1 EF-1-gamma C-terminal domain.,similarity:Contains 1 GST C-terminal domain.,similarity:Contains 1 GST N-terminal domain.,subunit:EF-1 is composed of four subunits: alpha, beta, delta, and gamma.,tissue specificity:Highly expressed in pancreatic tumor tissue and to a lesser extent in normal kidney, intestine, pancreas, stomach, lung, brain, spleen

Nanjing BYabscience technology Co.,Ltd







contains an N-terminal glutathione transferase domain, which may be involved in regulating the assembly of multisubunit complexes containing this elongation factor and aminoacyl-tRNA synthetases. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using EF-1 γ Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd