



ERF Monoclonal Antibody

Catalog No BYmab-01701 Isotype IgG Reactivity Human;Mouse Applications WB Gene Name ERF Protein Name ETS domain-containing transcription factor ERF Immunogen The antiserum was produced against synthesized peptide derived from human ERF. AA range:492-541 Specificity ERF Monoclonal Antibody detects endogenous levels of ERF protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms ERF; ETS domain-containing transcription factor ERF; Ets2 repressor factor; PE-2 Observed Band Cell Pathway Nucleus. Tissue Specificity Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required		
Reactivity Human; Mouse Applications WB Gene Name ERF Protein Name ETS domain-containing transcription factor ERF Immunogen The antiserum was produced against synthesized peptide derived from human ERF. AA range: 492-541 Specificity ERF Monoclonal Antibody detects endogenous levels of ERF protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms ERF; ETS domain-containing transcription factor ERF; Ets2 repressor factor; PE-2 Observed Band Cell Pathway Nucleus. Tissue Specificity Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and chorioallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation, errophorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF. sequence caution: The sequence differs from that shown because it seems to be derived from a pre-mRNA, similarity. Belongs to the ETS family, similarity. Contains 1 ETS DNA-binding domain, tissue specificity: Highest	Catalog No	BYmab-01701
Applications Gene Name ERF Protein Name ETS domain-containing transcription factor ERF Immunogen The antiserum was produced against synthesized peptide derived from human ERF. AA range: 492-541 Specificity ERF Monoclonal Antibody detects endogenous levels of ERF protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms ERF; ETS domain-containing transcription factor ERF; Ets2 repressor factor; PE-2 Observed Band Cell Pathway Nucleus. Tissue Specificity Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and choricoallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation, errophorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF. sequence caution: The sequence differs from that shown because it seems to be derived from a pre-mRNA, similarity. Belongs to the ETS family, similarity; Contains 1 ETS DNA-binding domain, tissue specificity: Highest	Isotype	IgG
Gene Name ETS domain-containing transcription factor ERF Immunogen The antiserum was produced against synthesized peptide derived from human ERF. AA range:492-541 Specificity ERF Monoclonal Antibody detects endogenous levels of ERF protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms ERF; ETS domain-containing transcription factor ERF; Ets2 repressor factor; PE-2 Observed Band Cell Pathway Nucleus. Tissue Specificity Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and choricallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation., PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF; Sequence califers from that show because it seems to be derived from a pre-mRNA, similarity:Belongs to the ETS family, similarity:Colonians 1 ETS DNA-binding domain., itsue specificity:Highest	Reactivity	Human;Mouse
Protein Name Immunogen The antiserum was produced against synthesized peptide derived from human ERF. AA range:492-541 Specificity ERF Monoclonal Antibody detects endogenous levels of ERF protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms ERF; ETS domain-containing transcription factor ERF; Ets2 repressor factor; PE-2 Observed Band Cell Pathway Nucleus. Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, etoplacental cone cavity closure, and choricalalantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation., PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF, sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA, similarity:Belongs to the ETS family, similarity:Contains 1 ETS DNA-binding domain., tissue specificity:Highest	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human ERF. AA range:492-541 Specificity ERF Monoclonal Antibody detects endogenous levels of ERF protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml 290% Storage Stability -20°C/1 year Synonyms ERF; ETS domain-containing transcription factor ERF; Ets2 repressor factor; PE-2 Observed Band Cell Pathway Nucleus. Tissue Specificity Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and choricallantic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation, PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF, sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA, similarity. Belongs to the ETS family, similarity. Contains 1 ETS DNA-binding domain., tissue specificity:Highest	Gene Name	ERF
ERF. AA range:492-541 Specificity ERF Monoclonal Antibody detects endogenous levels of ERF protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms ERF; ETS domain-containing transcription factor ERF; Ets2 repressor factor; PE-2 Observed Band Cell Pathway Nucleus. Tissue Specificity Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and chorioallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation, PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF., sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA, similarity. Belongs to the ETS family, similarity. Contains 1 ETS DNA-binding domain., tissue specificity:Highest	Protein Name	ETS domain-containing transcription factor ERF
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms ERF; ETS domain-containing transcription factor ERF; Ets2 repressor factor; PE-2 Observed Band Cell Pathway Nucleus. Tissue Specificity Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and chorioallantoic attachment (By similanty). May be important for regulating trophoblast stem cell differentiation. PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylates the activity of ERF., sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA, similarity:Belongs to the ETS family, similarity:Contains 1 ETS DNA-binding domain, tissue specificity:Highest	Immunogen	
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms ERF; ETS domain-containing transcription factor ERF; Ets2 repressor factor; PE-2 Observed Band Cell Pathway Nucleus. Tissue Specificity Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and chorioallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation., PTM:Phosphorylated by multiple kinases including probable pERK2. Phosphorylated by multiple kinases including probable pERK2. Phosphorylation regulates the activity of ERF., sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA., similarity:Belongs to the ETS family., similarity:Contains 1 ETS DNA-binding domain., tissue specificity:Highest	Specificity	ERF Monoclonal Antibody detects endogenous levels of ERF protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms ERF; ETS domain-containing transcription factor ERF; Ets2 repressor factor; PE-2 Observed Band Cell Pathway Nucleus. Tissue Specificity Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectopacental cone cavity closure, and chorioallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation.,PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF.,sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA, similarity:Belongs to the ETS family, similarity:Contains 1 ETS DNA-binding domain, tissue specificity:Highest	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms ERF; ETS domain-containing transcription factor ERF; Ets2 repressor factor; PE-2 Observed Band Cell Pathway Nucleus. Tissue Specificity Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and chorioallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation., PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF., sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA, similarity:Belongs to the ETS family.,similarity:Contains 1 ETS DNA-binding domain, itissue specificity:Highest	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms ERF; ETS domain-containing transcription factor ERF; Ets2 repressor factor; PE-2 Observed Band Cell Pathway Nucleus. Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and chorioallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation., PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF., sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA, similarity:Belongs to the ETS family, similarity:Contains 1 ETS DNA-binding domain., tissue specificity:Highest	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms ERF; ETS domain-containing transcription factor ERF; Ets2 repressor factor; PE-2 Observed Band Cell Pathway Nucleus. Tissue Specificity Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and chorioallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation.,PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF.,sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA.,similarity:Belongs to the ETS family.,similarity:Contains 1 ETS DNA-binding domain.,tissue specificity:Highest	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year ERF; ETS domain-containing transcription factor ERF; Ets2 repressor factor; PE-2 Observed Band Cell Pathway Nucleus. Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and chorioallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation.,PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF.,sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA.similarity:Belongs to the ETS family.,similarity:Contains 1 ETS DNA-binding domain.,tissue specificity:Highest	Concentration	1 mg/ml
Synonyms ERF; ETS domain-containing transcription factor ERF; Ets2 repressor factor; PE-2 Observed Band Cell Pathway Nucleus. Tissue Specificity Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and chorioallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation.,PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF.,sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA.,similarity:Belongs to the ETS family.,similarity:Contains 1 ETS DNA-binding domain.,tissue specificity:Highest	Purity	≥90%
Observed Band Cell Pathway Nucleus. Tissue Specificity Highest levels in testis, ovary, pancreas, and heart. Function function:Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and chorioallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation.,PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF.,sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA.,similarity:Belongs to the ETS family.,similarity:Contains 1 ETS DNA-binding domain.,tissue specificity:Highest	Storage Stability	-20°C/1 year
Cell Pathway Nucleus. Tissue Specificity Highest levels in testis, ovary, pancreas, and heart. function: Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and chorioallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation.,PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF.,sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA.,similarity:Belongs to the ETS family.,similarity:Contains 1 ETS DNA-binding domain.,tissue specificity:Highest	Synonyms	·
Tissue Specificity Highest levels in testis, ovary, pancreas, and heart. function: Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and chorioallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation.,PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF.,sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA.,similarity:Belongs to the ETS family.,similarity:Contains 1 ETS DNA-binding domain.,tissue specificity:Highest	Observed Band	
function: Potent transcriptional repressor that binds to the H1 element of the Ets2 promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and chorioallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation.,PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF.,sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA.,similarity:Belongs to the ETS family.,similarity:Contains 1 ETS DNA-binding domain.,tissue specificity:Highest	Cell Pathway	Nucleus.
promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and chorioallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation.,PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF.,sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA.,similarity:Belongs to the ETS family.,similarity:Contains 1 ETS DNA-binding domain.,tissue specificity:Highest	Tissue Specificity	Highest levels in testis, ovary, pancreas, and heart.
	Function	promoter. May regulate other genes involved in cellular proliferation. Required for extraembryonic ectoderm differentiation, ectoplacental cone cavity closure, and chorioallantoic attachment (By similarity). May be important for regulating trophoblast stem cell differentiation.,PTM:Phosphorylated by multiple kinases including probably ERK2. Phosphorylation regulates the activity of ERF.,sequence caution:The sequence differs from that shown because it seems to be derived from a pre-mRNA.,similarity:Belongs to the ETS family.,similarity:Contains 1 ETS DNA-binding domain.,tissue specificity:Highest

Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background	ETS2 is a transcription factor and protooncogene involved in development, apoptosis, and the regulation of telomerase. The protein encoded by this gene binds to the ETS2 promoter and is a strong repressor of ETS2 transcription. Several transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Aug 2015],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using ERF Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd