



FoxN4 Monoclonal Antibody

Catalog No BYmab-01731 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name FOXN4 Protein Name Forkhead box protein N4 Immunogen The antiserum was produced against synthesized peptide derived from human FOXN4. AA range:411-460 Specificity FoxN4 Monoclonal Antibody detects endogenous levels of FoxN4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FOXN4; Forkhead box protein N4 Observed Band 50kD Cell Pathway Nucleus Tissue Specificity Brain, Clones donated by RIKEN, Function similarity: Contains 1 fork-head DNA-binding domain. Background Members of the winged-helix/forkhead family of transcription factors are involved in a v		
Reactivity Human;Rat;Mouse; Applications WB Gene Name FOXN4 Protein Name Forkhead box protein N4 Immunogen The antiserum was produced against synthesized peptide derived from human FOXN4. AA range:411-460 Specificity FoxN4 Monoclonal Antibody detects endogenous levels of FoxN4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FOXN4; Forkhead box protein N4 Observed Band 50kD Cell Pathway Nucleus . Tissue Specificity Brain, Clones donated by RIKEN, Function similarity:Contains 1 fork-head DNA-binding domain., Members of the winged-helix/forkhead family of transcription factors, such as FOXN4; are characterized by a 110-amino acid DNA-binding domain that can fold into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes a key regulators in development and metabolism (Li et al.,	Catalog No	BYmab-01731
Applications Gene Name FOXN4 Protein Name Forkhead box protein N4 Immunogen The antiserum was produced against synthesized peptide derived from human FOXN4. AA range:411-460 Specificity FoxN4 Monoclonal Antibody detects endogenous levels of FoxN4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FOXN4; Forkhead box protein N4 Observed Band 50kD Cell Pathway Nucleus. Tissue Specificity Brain, Clones donated by RIKEN, Function Similarity: Contains 1 fork-head DNA-binding domain., Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a varient of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and marked by a final constraint of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Isotype	IgG
Gene Name FOXN4 Protein Name Forkhead box protein N4 Immunogen The antiserum was produced against synthesized peptide derived from human FOXN4. AA range:411-460 Specificity FoxN4 Monoclonal Antibody detects endogenous levels of FoxN4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FOXN4; Forkhead box protein N4 Observed Band 50kD Cell Pathway Nucleus. Tissue Specificity Brain, Clones donated by RIKEN, Function similarity: Contains 1 fork-head DNA-binding domain., Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a varient of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Reactivity	Human;Rat;Mouse;
Protein Name Forkhead box protein N4 Immunogen The antiserum was produced against synthesized peptide derived from human FOXN4. AA range:411-460 Specificity FoxN4 Monoclonal Antibody detects endogenous levels of FoxN4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FOXN4; Forkhead box protein N4 Observed Band 50kD Cell Pathway Nucleus . Tissue Specificity Brain,Clones donated by RIKEN, Function similarity:Contains 1 fork-head DNA-binding domain., Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and eatelolism (Li et al.,	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human FOXN4. AA range:411-460 Specificity FoxN4 Monoclonal Antibody detects endogenous levels of FoxN4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms FOXN4; Forkhead box protein N4 Observed Band 50kD Cell Pathway Nucleus Tissue Specificity Brain, Clones donated by RIKEN, Function similarity:Contains 1 fork-head DNA-binding domain., Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by a large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Gene Name	FOXN4
FOXN4. AA range:411-460 Specificity FoxN4 Monoclonal Antibody detects endogenous levels of FoxN4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FOXN4; Forkhead box protein N4 Observed Band 50kD Cell Pathway Nucleus Tissue Specificity Brain, Clones donated by RIKEN, Function Background Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Protein Name	Forkhead box protein N4
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FOXN4; Forkhead box protein N4 Observed Band 50kD Cell Pathway Nucleus Tissue Specificity Brain,Clones donated by RIKEN, Function Similarity:Contains 1 fork-head DNA-binding domain. Background Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key transcription in development and metabolism (Li et al.,	Immunogen	
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FOXN4; Forkhead box protein N4 Observed Band 50kD Cell Pathway Nucleus . Tissue Specificity Brain, Clones donated by RIKEN, Function similarity: Contains 1 fork-head DNA-binding domain., Background Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Specificity	FoxN4 Monoclonal Antibody detects endogenous levels of FoxN4 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FOXN4; Forkhead box protein N4 Observed Band 50kD Cell Pathway Nucleus . Tissue Specificity Brain,Clones donated by RIKEN, Function Similarity:Contains 1 fork-head DNA-binding domain., Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FOXN4; Forkhead box protein N4 Observed Band 50kD Cell Pathway Nucleus . Tissue Specificity Brain, Clones donated by RIKEN, Function similarity:Contains 1 fork-head DNA-binding domain., Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FOXN4; Forkhead box protein N4 Observed Band 50kD Cell Pathway Nucleus . Tissue Specificity Brain,Clones donated by RIKEN, Function similarity:Contains 1 fork-head DNA-binding domain., Background Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms FOXN4; Forkhead box protein N4 Observed Band 50kD Cell Pathway Nucleus . Tissue Specificity Brain, Clones donated by RIKEN, Function similarity: Contains 1 fork-head DNA-binding domain., Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can folcinto a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year FOXN4; Forkhead box protein N4 Observed Band 50kD Cell Pathway Nucleus . Tissue Specificity Brain, Clones donated by RIKEN, Function Similarity: Contains 1 fork-head DNA-binding domain., Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Concentration	1 mg/ml
Synonyms FOXN4; Forkhead box protein N4 Observed Band 50kD Cell Pathway Nucleus . Tissue Specificity Brain, Clones donated by RIKEN, Function similarity: Contains 1 fork-head DNA-binding domain., Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Purity	≥90%
Observed Band Cell Pathway Nucleus . Tissue Specificity Brain, Clones donated by RIKEN, Function Similarity: Contains 1 fork-head DNA-binding domain., Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Storage Stability	-20°C/1 year
Cell Pathway Nucleus . Tissue Specificity Brain,Clones donated by RIKEN, Function similarity:Contains 1 fork-head DNA-binding domain., Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Synonyms	FOXN4; Forkhead box protein N4
Tissue Specificity Brain, Clones donated by RIKEN, similarity: Contains 1 fork-head DNA-binding domain., Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Observed Band	50kD
Function Similarity:Contains 1 fork-head DNA-binding domain., Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can folcinto a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Cell Pathway	Nucleus .
Background Members of the winged-helix/forkhead family of transcription factors, such as FOXN4, are characterized by a 110-amino acid DNA-binding domain that can folconinto a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Tissue Specificity	Brain,Clones donated by RIKEN,
FOXN4, are characterized by a 110-amino acid DNA-binding domain that can folcond into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,	Function	similarity:Contains 1 fork-head DNA-binding domain.,
	Background	FOXN4, are characterized by a 110-amino acid DNA-binding domain that can fold into a variant of the helix-turn-helix motif consisting of 3 alpha helices flanked by 2 large loops or wings. These transcription factors are involved in a variety of biologic processes as key regulators in development and metabolism (Li et al.,

Nanjing BYabscience technology Co.,Ltd






matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using FoxN4 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd