



# HRT2 Monoclonal Antibody

Catalog No	BYmab-01815
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	HEY2
Protein Name	Hairy/enhancer-of-split related with YRPW motif protein 2
Immunogen	The antiserum was produced against synthesized peptide derived from human HEY2. AA range:21-70
Specificity	HRT2 Monoclonal Antibody detects endogenous levels of HRT2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HEY2; BHLHB32; CHF1; GRL; HERP; HERP1; HRT2; Hairy/enhancer-of-split related with YRPW motif protein 2; Cardiovascular helix-loop-helix factor 1; hCHF1; Class B basic helix-loop-helix protein 32; bHLHb32; HES-related repressor protein 2; Ha
Observed Band	36kD
Cell Pathway	Nucleus .
Tissue Specificity	Heart,Lung,Testis,
Function	disease:Defects in HEY2 may be involved in atrioventricular septal defects (AVSD).,function:Downstream effector of Notch signaling which may be required for cardiovascular development. Transcriptional repressor which binds preferentially to the canonical E box sequence 5'-CACGTG-3'. Represses transcription by the cardiac transcriptional activators GATA4 and GATA6.,similarity:Belongs to the HEY family.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,similarity:Contains 1 Orange

Nanjing BYabscience technology Co.,Ltd



domain.,subunit:May self-associate (By similarity). Interacts with GATA4, HES1 and HEYL (By similarity). Interacts with HDAC1, NCOR1 and SIN3A (By similarity). Interacts with ARNT and GATA6.,

#### Background

This gene encodes a member of the hairy and enhancer of split-related (HESR) family of basic helix-loop-helix (bHLH)-type transcription factors. The encoded protein forms homo- or hetero-dimers that localize to the nucleus and interact with a histone deacetylase complex to repress transcription. Expression of this gene is induced by the Notch signal transduction pathway. Two similar and redundant genes in mouse are required for embryonic cardiovascular development, and are also implicated in neurogenesis and somitogenesis. Alternatively spliced transcript variants have been found, but their biological validity has not been determined. [provided by RefSeq, Jul 2008],


#### matters needing attention

Avoid repeated freezing and thawing!

#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

### Products Images



Nanjing BYabscience technology Co.,Ltd