



# NAB2 Monoclonal Antibody

Catalog No	BYmab-01889
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	NAB2
Protein Name	NGFI-A-binding protein 2
Immunogen	The antiserum was produced against synthesized peptide derived from human NAB2. AA range:261-310
Specificity	NAB2 Monoclonal Antibody detects endogenous levels of NAB2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NAB2; MADER; NGFI-A-binding protein 2; EGR-1-binding protein 2; Melanoma-associated delayed early response protein; Protein MADER
Observed Band	56kD
Cell Pathway	Nucleus . Isoform 2 is not localized to the nucleus. .
Tissue Specificity	Widely expressed at low levels. Highly expressed in melanoma cell lines.
Function	domain:The NAB conserved domain 1 (NCD1) interacts with EGR1 inhibitory domain and mediates multimerization.,domain:The NAB conserved domain 2 (NCD2) is necessary for transcriptional repression.,function:Acts as a transcriptional repressor for zinc finger transcription factors EGR1 and EGR2. Isoform 2 lacks repression ability.,induction:By serum and PMA stimulation.,similarity:Belongs to the NAB family.,subcellular location:Isoform 2 is not localized to the nucleus.,subunit:Homomultimers may associate with EGR1 bound to DNA.,tissue specificity:Widely expressed at low levels. Highly expressed in melanoma cell lines.,

Nanjing BYabscience technology Co.,Ltd



## Background

This gene encodes a member of the family of NGFI-A binding (NAB) proteins, which function in the nucleus to repress transcription induced by some members of the EGR (early growth response) family of transactivators. NAB proteins can homo- or hetero-multimerize with other EGR or NAB proteins through a conserved N-terminal domain, and repress transcription through two partially redundant C-terminal domains. Transcriptional repression by the encoded protein is mediated in part by interactions with the nucleosome remodeling and deacetylase (NuRD) complex. Alternatively spliced transcript variants have been described, but their biological validity has not been determined. [provided by RefSeq, Jul 2008],


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using NAB2 Monoclonal Antibody