



Nap1 Monoclonal Antibody

Catalog No	BYmab-01891
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	AZI2
Protein Name	5-azacytidine-induced protein 2
Immunogen	The antiserum was produced against synthesized peptide derived from human AZI2. AA range:10-59
Specificity	Nap1 Monoclonal Antibody detects endogenous levels of Nap1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year AZI2; NAP1; 5-azacytidine-induced protein 2; NF-kappa-B-activating kinase-associated protein 1; Nak-associated protein 1; TILP
	AZI2; NAP1; 5-azacytidine-induced protein 2; NF-kappa-B-activating
Synonyms	AZI2; NAP1; 5-azacytidine-induced protein 2; NF-kappa-B-activating kinase-associated protein 1; Nak-associated protein 1; TILP
Synonyms Observed Band	AZI2; NAP1; 5-azacytidine-induced protein 2; NF-kappa-B-activating kinase-associated protein 1; Nak-associated protein 1; TILP 48kD
Synonyms Observed Band Cell Pathway	AZI2; NAP1; 5-azacytidine-induced protein 2; NF-kappa-B-activating kinase-associated protein 1; Nak-associated protein 1; TILP 48kD Cytoplasm . Widely expressed (PubMed:14560022). Abundant expression seen in the pancreas and testis (PubMed:14560022). function:Activates serine/threonine-protein kinase TBK1 and facilitates its
Synonyms Observed Band Cell Pathway Tissue Specificity	AZI2; NAP1; 5-azacytidine-induced protein 2; NF-kappa-B-activating kinase-associated protein 1; Nak-associated protein 1; TILP 48kD Cytoplasm. Widely expressed (PubMed:14560022). Abundant expression seen in the pancreas and testis (PubMed:14560022). function:Activates serine/threonine-protein kinase TBK1 and facilitates its oligomerization. Enhances the phosphorylation of NF-kappa-B p65 subunit RELA by TBK1. Promotes TBK1-induced as well as TNF-alpha or PMA-induced activation of NF-kappa-B. Participates in IFN-beta promoter activation via TICAM1.,subunit:Interacts with IKBKE, TBK1 and TICAM1.,tissue specificity:Widely expressed. Abundant expression seen in the pancreas and


Nanjing BYabscience technology Co.,Ltd



matters needing

Usage suggestions


attention





NAK (TBK1; MIM 604834) (Fujita et al., 2003 [PubMed 14560022]).[supplied by OMIM, Mar 2008],	
Avoid repeated freezing and thawing!	
This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	

Products Images



Western Blot analysis of various cells using Nap1 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd