



POLDIP3 Monoclonal Antibody

with POLD2.,* Background This gene encodes an RRM (RNA recognition motif)-containing protein that		
Reactivity Human;Rat;Mouse; Applications WB Gene Name POLDIP3 Protein Name Polymerase delta-interacting protein 3 Immunogen The antiserum was produced against synthesized peptide derived from human POLDIP3. AA range:348-397 Specificity POLDIP3 Monoclonal Antibody detects endogenous levels of POLDIP3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms POLDIP3; KIAA1649; PDIP46; Polymerase delta-interacting protein 3; 46 kDa DNA polymerase delta interaction protein; p46; S6K1 Aly/REF-like target; SKAR Observed Band 46kD Cell Pathway Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain,Epithelium,Ovarian carcinoma,Placenta,Skin,Testis, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity:Contains 1 RRM (RNA recognition motif) domain., subunit:Interacts with POLD2., Background This gene encodes an RRM (RNA recognition motif)-containing protein fa6 kinase beta-1 to mRNAs. Alternative splicing results in multipte rainsoript variants.	Catalog No	BYmab-01945
Applications WB Gene Name POLDIP3 Protein Name Polymerase delta-interacting protein 3 Immunogen The antiserum was produced against synthesized peptide derived from human POLDIP3. AA range:348-397 Specificity POLDIP3 Monoclonal Antibody detects endogenous levels of POLDIP3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms POLDIP3; KIAA1649; PDIP46; Polymerase delta-interacting protein 3; 46 kDa DNA polymerase delta interaction protein; p46; S6K1 Aly/REF-like target; SKAR Observed Band 46kD Cell Pathway Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain,Epithelium,Ovarian carcinoma,Placenta,Skin,Testis, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,simillarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Interacts with POLD2. Background This gene encodes an RRM (RNA recognition motif) protein that participates in the regulation of translation by recruiting ribosomal protein Staring protein that participates in the regulation of translation by recruiting ribosomal protein Staring protein that participates in the regulation of translation by recruiting ribosomal protein Staring protein that participates in the regulation of translation by recruiting ribosomal protein Staring Protein St	Isotype	IgG
Gene Name POLDIP3 Protein Name Polymerase delta-interacting protein 3 Immunogen The antiserum was produced against synthesized peptide derived from human POLDIP3. AA range:348-397 Specificity POLDIP3 Monoclonal Antibody detects endogenous levels of POLDIP3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms POLDIP3; KIAA1649; PDIP46; Polymerase delta-interacting protein 3; 46 kDa DNA polymerase delta interaction protein; p46; S6K1 Aly/REF-like target; SKAR Observed Band 46kD Cell Pathway Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain, Epithelium, Ovarian carcinoma, Placenta, Skin, Testis, Function PTM: Phosphorylated upon DNA damage, probably by ATM or ATR, similarity-Contains 1 RRM (RNA recognition motif) domain, subunit: Interacts with POLD2 Background This gene encodes an RRM (RNA recognition	Reactivity	Human;Rat;Mouse;
Protein Name Polymerase delta-interacting protein 3 Immunogen The antiserum was produced against synthesized peptide derived from human POLDIP3. AA range:348-397 Specificity POLDIP3 Monoclonal Antibody detects endogenous levels of POLDIP3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms POLDIP3; KIAA1649; PDIP46; Polymerase delta-interacting protein 3; 46 kDa DNA polymerase delta interaction protein; p46; S6K1 Aly/REF-like target; SKAR Observed Band 46kD Cell Pathway Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain,Epithelium,Ovarian carcinoma,Placenta,Skin,Testis, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR, similarity. Contains 1 RRM (RNA recognition motif) domain., subunit:Interacts with POLD2., Background This gene encodes an RRM (RNA recognition motif) containing protein that participates in the regulation of translati	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human POLDIP3. AA range:348-397 Specificity POLDIP3 Monoclonal Antibody detects endogenous levels of POLDIP3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms POLDIP3; KIAA1649; PDIP46; Polymerase delta-interacting protein 3; 46 kDa DNA polymerase delta interaction protein; p46; S6K1 Aly/REF-like target; SKAR Observed Band 46kD Cell Pathway Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain,Epithelium,Ovarian carcinoma,Placenta,Skin,Testis, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR. similarity:Contains 1 RRM (RNA recognition motif) domain., subunit:Interacts with POLD2. Background This gene encodes an RRM (RNA recognition motif)-containing protein that participates in the regulation of translation by recruiting inbosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple trans	Gene Name	POLDIP3
POLDIP3. AA range:348-397 Specificity POLDIP3 Monoclonal Antibody detects endogenous levels of POLDIP3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms POLDIP3; KIAA1649; PDIP46; Polymerase delta-interacting protein 3; 46 kDa DNA polymerase delta interaction protein; p46; S6K1 Aly/REF-like target; SKAR Observed Band 46kD Cell Pathway Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain,Epithelium,Ovarian carcinoma,Placenta,Skin,Testis, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.;similarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Interacts with POLD2 Background This gene encodes an RRM (RNA recognition motif)-containing protein that participates in the regulation of translation by recruiting ibosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.	Protein Name	Polymerase delta-interacting protein 3
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms POLDIP3; KIAA1649; PDIP46; Polymerase delta-interacting protein 3; 46 kDa DNA polymerase delta interaction protein; p46; S6K1 Aly/REF-like target; SKAR Observed Band 46kD Cell Pathway Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain,Epithelium,Ovarian carcinoma,Placenta,Skin,Testis, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR. similarity:Contains 1 RRM (RNA recognition motif) domainsubunit:Interacts with POLD2., With POLD2. This gene encodes an RRM (RNA recognition motif)-containing protein that participates in the regulation of translation by recruiting ribosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.	Immunogen	
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms POLDIP3; KIAA1649; PDIP46; Polymerase delta-interacting protein 3; 46 kDa DNA polymerase delta interaction protein; p46; S6K1 Aly/REF-like target; SKAR Observed Band 46kD Cell Pathway Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain,Epithelium,Ovarian carcinoma,Placenta,Skin,Testis, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Interacts with POLD2., Background This gene encodes an RRM (RNA recognition motif)-containing protein 56 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.	Specificity	POLDIP3 Monoclonal Antibody detects endogenous levels of POLDIP3 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms POLDIP3; KIAA1649; PDIP46; Polymerase delta-interacting protein 3; 46 kDa DNA polymerase delta interaction protein; p46; S6K1 Aly/REF-like target; SKAR Observed Band 46kD Cell Pathway Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain,Epithelium,Ovarian carcinoma,Placenta,Skin,Testis, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR, similiarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Interacts with POLD2., Background This gene encodes an RRM (RNA recognition motif)-containing protein that participates in the regulation of translation by recruiting ribosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms POLDIP3; KIAA1649; PDIP46; Polymerase delta-interacting protein 3; 46 kDa DNA polymerase delta interaction protein; p46; S6K1 Aly/REF-like target; SKAR Observed Band 46kD Cell Pathway Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain,Epithelium,Ovarian carcinoma,Placenta,Skin,Testis, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Interacts with POLD2., Background This gene encodes an RRM (RNA recognition motif)-containing protein that participates in the regulation of translation by recruiting ribosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms POLDIP3; KIAA1649; PDIP46; Polymerase delta-interacting protein 3; 46 kDa DNA polymerase delta interaction protein; p46; S6K1 Aly/REF-like target; SKAR Observed Band 46kD Cell Pathway Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain,Epithelium,Ovarian carcinoma,Placenta,Skin,Testis, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Interacts with POLD2., Background This gene encodes an RRM (RNA recognition motif)-containing protein that participates in the regulation of translation by recruiting ribosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.	Purification	•
Purity ≥90% Storage Stability -20°C/1 year Synonyms POLDIP3; KIAA1649; PDIP46; Polymerase delta-interacting protein 3; 46 kDa DNA polymerase delta interaction protein; p46; S6K1 Aly/REF-like target; SKAR Observed Band 46kD Cell Pathway Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain,Epithelium,Ovarian carcinoma,Placenta,Skin,Testis, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Interacts with POLD2., Background This gene encodes an RRM (RNA recognition motif)-containing protein that participates in the regulation of translation by recruiting ribosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms POLDIP3; KIAA1649; PDIP46; Polymerase delta-interacting protein 3; 46 kDa DNA polymerase delta interaction protein; p46; S6K1 Aly/REF-like target; SKAR Observed Band 46kD Cell Pathway Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain,Epithelium,Ovarian carcinoma,Placenta,Skin,Testis, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Interacts with POLD2., Background This gene encodes an RRM (RNA recognition motif)-containing protein that participates in the regulation of translation by recruiting ribosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.	Concentration	1 mg/ml
Synonyms POLDIP3; KIAA1649; PDIP46; Polymerase delta-interacting protein 3; 46 kDa DNA polymerase delta interaction protein; p46; S6K1 Aly/REF-like target; SKAR Observed Band 46kD Cell Pathway Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain,Epithelium,Ovarian carcinoma,Placenta,Skin,Testis, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Interacts with POLD2., Background This gene encodes an RRM (RNA recognition motif)-containing protein that participates in the regulation of translation by recruiting ribosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.	Purity	≥90%
DNA polymerase delta interaction protein; p46; S6K1 Aly/REF-like target; SKAR Observed Band 46kD Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain,Epithelium,Ovarian carcinoma,Placenta,Skin,Testis, Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Interacts with POLD2., Background This gene encodes an RRM (RNA recognition motif)-containing protein that participates in the regulation of translation by recruiting ribosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.	Storage Stability	-20°C/1 year
Cell Pathway Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein. Tissue Specificity Brain, Epithelium, Ovarian carcinoma, Placenta, Skin, Testis, Function PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Contains 1 RRM (RNA recognition motif) domain., subunit: Interacts with POLD2., Background This gene encodes an RRM (RNA recognition motif)-containing protein that participates in the regulation of translation by recruiting ribosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.	Synonyms	
Tissue Specificity Brain, Epithelium, Ovarian carcinoma, Placenta, Skin, Testis, PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Contains 1 RRM (RNA recognition motif) domain., subunit: Interacts with POLD2., Background This gene encodes an RRM (RNA recognition motif)-containing protein that participates in the regulation of translation by recruiting ribosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.	Observed Band	46kD
Function PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Interacts with POLD2., Background This gene encodes an RRM (RNA recognition motif)-containing protein that participates in the regulation of translation by recruiting ribosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.	Cell Pathway	Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein.
ATR.,similarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Interacts with POLD2., Background This gene encodes an RRM (RNA recognition motif)-containing protein that participates in the regulation of translation by recruiting ribosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.	Tissue Specificity	Brain, Epithelium, Ovarian carcinoma, Placenta, Skin, Testis,
participates in the regulation of translation by recruiting ribosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.	Function	ATR.,similarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Interacts
	Background	participates in the regulation of translation by recruiting ribosomal protein S6 kinase beta-1 to mRNAs. Alternative splicing results in multiple transcript variants.

Nanjing BYabscience technology Co.,Ltd





matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

