



# RMP Monoclonal Antibody

<b>Catalog No</b>	BYmab-01983
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	URI1
<b>Protein Name</b>	Unconventional prefoldin RPB5 interactor 1
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human RMP. AA range:415-464
<b>Specificity</b>	RMP Monoclonal Antibody detects endogenous levels of RMP protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	URI1; C19orf2; NNX3; PPP1R19; RMP; URI; Unconventional prefoldin RPB5 interactor 1; Protein NNX3; Protein phosphatase 1 regulatory subunit 19; RNA polymerase II subunit 5-mediating protein; RPB5-mediating protein
<b>Observed Band</b>	56kD
<b>Cell Pathway</b>	Nucleus. Cytoplasm. Mitochondrion. Cell projection, dendrite . Colocalizes with PFDN2, PFDN4, PPP1CC, RPS6KB1 and STAP1 at mitochondrion.
<b>Tissue Specificity</b>	Ubiquitous. Expressed in ovarian cancers (at protein level). Expressed strongly in skeletal muscle. Expressed weakly in brain, heart, pancreas and in prostate epithelial cells.
<b>Function</b>	function:Seems to be a scaffolding protein able to assemble a prefoldin-like complex that contains PFDs and proteins with roles in transcription and ubiquitination.,similarity:Belongs to the RNA polymerase II subunit 5-mediating protein family.,subunit:Component of the URI complex that contains PFD2, POLR2E/RPB5, TIP48, TIP49 and URI. Interacts with POLR2E/RPB5 and LRPPRC.,tissue specificity:Ubiquitous.,

**Nanjing BYabscience technology Co.,Ltd**



## Background

This gene encodes member of the prefoldin family of molecular chaperones. The encoded protein functions as a scaffolding protein and plays roles in ubiquitination and transcription, in part through interactions with the RNA polymerase II subunit RPB5. This gene may play a role in multiple malignancies including ovarian cancer and hepatocellular carcinoma. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and a pseudogene of this gene is located on the long arm of chromosome 22. [provided by RefSeq, Nov 2011],


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using RMP Monoclonal Antibody

nce technology Co.,Ltd