

SRp75 Monoclonal Antibody

Catalog No BYmab-02047 Isotype IgG Reactivity Human;Mouse Applications WB Gene Name SRSF4 Protein Name Serine/arginine-rich splicing factor 4 Immunogen The antiserum was produced against synthesized peptide derived from human SFRS4. AA range:111-160 Specificity SRp75 Monoclonal Antibody detects endogenous levels of SRp75 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SRSF4; SFRS4; SRP75; Serine/arginine-rich splicing factor 4; Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, Function function: A probable role in alternative splice site s		
Reactivity Human; Mouse Applications WB Gene Name SRSF4 Protein Name Serine/arginine-rich splicing factor 4 Immunogen The antiserum was produced against synthesized peptide derived from human SFRS4. AA range:111-160 Specificity SRp75 Monoclonal Antibody detects endogenous levels of SRp75 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SRSF4; SFRS4; SRP75; Serine/arginine-rich splicing factor 4; Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle . Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, function: A probable role in alternative splice site selection during pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN., Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Catalog No	BYmab-02047
Applications Gene Name SRSF4 Protein Name Serine/arginine-rich splicing factor 4 Immunogen The antiserum was produced against synthesized peptide derived from human SFRS4. AA range:111-160 Specificity SRp75 Monoclonal Antibody detects endogenous levels of SRp75 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SRSF4; SFRS4; SRP75; Serine/arginine-rich splicing factor 4; Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, function: A probable role in alternative splice site selection during pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN., Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Isotype	IgG
Gene Name Protein Name Serine/arginine-rich splicing factor 4 Immunogen The antiserum was produced against synthesized peptide derived from human SFRS4. AA range:111-160 Specificity SRp75 Monoclonal Antibody detects endogenous levels of SRp75 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SRSF4; SFRS4; SRP75; Serine/arginine-rich splicing factor 4; Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle. Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, function: A probable role in alternative splice site selection during pre-mRNA splicing, PTM: Extensively phosphorylated on serine residues in the RS domain, similarity: Belongs to the splicing factor SR family, similarity. Contains 2 cRRM (RNA recognition motif) domains, subunit-Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN. Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Reactivity	Human;Mouse
Protein Name Serine/arginine-rich splicing factor 4 Immunogen The antiserum was produced against synthesized peptide derived from human SFRS4. AA range:111-160 Specificity SRp75 Monoclonal Antibody detects endogenous levels of SRp75 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SRSF4; SFRS4; SRP75; Serine/arginine-rich splicing factor 4; Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle . Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, Function function: A probable role in alternative splice site selection during pre-mRNA splicing, PTM: Extensively phosphorylated on serine residues in the RS domain, similarity: Belongs to the splicing factor SR family, similarity: Contains 2 RRM (RNA recognition motif) domains, subunit: Found a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN.,	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human SFRS4. AA range:111-160 Specificity SRp75 Monoclonal Antibody detects endogenous levels of SRp75 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SRSF4; SFRS4; SRP75; Serine/arginine-rich splicing factor 4; Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle . Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, Function function: A probable role in alternative splice site selection during pre-mRNA splicing, PTM: Extensively phosphorylated on serine residues in the RS domain, similarity. Centains 2 RRM (RNA recognition motif) domains, subunit: Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN., Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Gene Name	SRSF4
SFRS4. AA range:111-160 Specificity SRp75 Monoclonal Antibody detects endogenous levels of SRp75 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SRSF4; SFRS4; SRP75; Serine/arginine-rich splicing factor 4; Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle . Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, Function function: A probable role in alternative splice site selection during pre-mRNA splicing, PTM: Extensively phosphorylated on serine residues in the RS domain, similarity: Belongs to the splicing factor SR family, similarity: Contains 2 RRM (RNA recognition motif) domains, subunit: Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN. Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Protein Name	Serine/arginine-rich splicing factor 4
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SRSF4; SFRS4; SRP75; Serine/arginine-rich splicing factor 4; Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle . Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, Function function: A probable role in alternative splice site selection during pre-mRNA splicing. PTM: Extensively phosphorylated on serine residues in the RS domain. similarity: Belongs to the splicing factor family. similarity: Contains 2 RRM (RNA recognition motif) domains. subunit: Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN., Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Immunogen	
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SRSF4; SFRS4; SRP75; Serine/arginine-rich splicing factor 4; Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle . Tissue Specificity Aorta endothelial cell,Cerebrum,Epithelium,Lymph, Function function:A probable role in alternative splice site selection during pre-mRNA splicing. PTM:Extensively phosphorylated on serine residues in the RS domain . similarity.Belongs to the splicing factor SR family, similarity:Contains 2 RRM (RNA recognition motif) domains . subunit:Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN., This gene encodes a member of the arginine/serine-rich splicing factor family.	Specificity	SRp75 Monoclonal Antibody detects endogenous levels of SRp75 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SRSF4; SFRS4; SRP75; Serine/arginine-rich splicing factor 4; Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle. Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, Function function: A probable role in alternative splice site selection during pre-mRNA splicing, PTM: Extensively phosphorylated on serine residues in the RS domain, similarity: Belongs to the splicing factor SR family, similarity: Contains 2 RRM (RNA recognition motif) domains, subunit: Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN. Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SRSF4; SFRS4; SRP75; Serine/arginine-rich splicing factor 4; Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle . Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, Function function: A probable role in alternative splice site selection during pre-mRNA splicing, PTM:Extensively phosphorylated on serine residues in the RS domain., similarity: Belongs to the splicing factor SR family., similarity: Contains 2 RRM (RNA recognition motif) domains., subunit: Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN., This gene encodes a member of the arginine/serine-rich splicing factor family.	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms SRSF4; SFRS4; SRP75; Serine/arginine-rich splicing factor 4; Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle . Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, Function function: A probable role in alternative splice site selection during pre-mRNA splicing, PTM: Extensively phosphorylated on serine residues in the RS domain, similarity: Belongs to the splicing factor SR family., similarity: Contains 2 RRM (RNA recognition motif) domains., subunit: Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN., Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Purification	, , , , , , , , , , , , , , , , , , , ,
Purity ≥90% Storage Stability -20°C/1 year Synonyms SRSF4; SFRS4; SRP75; Serine/arginine-rich splicing factor 4; Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle . Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, Function function: A probable role in alternative splice site selection during pre-mRNA splicing., PTM:Extensively phosphorylated on serine residues in the RS domainsimilarity: Belongs to the splicing factor SR family., similarity: Contains 2 RRM (RNA recognition motif) domains., subunit: Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN., Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Dilution	WB 1:500-2000
Storage Stability Synonyms SRSF4; SFRS4; SRP75; Serine/arginine-rich splicing factor 4; Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle. Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, Function function: A probable role in alternative splice site selection during pre-mRNA splicing, PTM: Extensively phosphorylated on serine residues in the RS domain., similarity: Belongs to the splicing factor SR family., spimilarly: Contains 2 RRM (RNA recognition motif) domains, subunit: Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN., Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Concentration	1 mg/ml
Synonyms SRSF4; SFRS4; SRP75; Serine/arginine-rich splicing factor 4; Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle. Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, Function function: A probable role in alternative splice site selection during pre-mRNA splicing., PTM: Extensively phosphorylated on serine residues in the RS domain., similarity: Belongs to the splicing factor SR family., similarity: Contains 2 RRM (RNA recognition motif) domains., subunit: Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN., Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Purity	≥90%
Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor; arginine/serine-rich 4 Observed Band 57kD Cell Pathway Nucleus speckle. Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, Function function: A probable role in alternative splice site selection during pre-mRNA splicing., PTM: Extensively phosphorylated on serine residues in the RS domain., similarity: Belongs to the splicing factor SR family., similarity: Contains 2 RRM (RNA recognition motif) domains., subunit: Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN., Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Storage Stability	-20°C/1 year
Cell Pathway Nucleus speckle . Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, Function function: A probable role in alternative splice site selection during pre-mRNA splicing. PTM: Extensively phosphorylated on serine residues in the RS domain., similarity: Belongs to the splicing factor SR family., similarity: Contains 2 RRM (RNA recognition motif) domains., subunit: Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN., Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Synonyms	Pre-mRNA-splicing factor SRP75; SRP001LB; Splicing factor;
Tissue Specificity Aorta endothelial cell, Cerebrum, Epithelium, Lymph, function: A probable role in alternative splice site selection during pre-mRNA splicing., PTM: Extensively phosphorylated on serine residues in the RS domain., similarity: Belongs to the splicing factor SR family., similarity: Contains 2 RRM (RNA recognition motif) domains., subunit: Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN., Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Observed Band	57kD
Function function:A probable role in alternative splice site selection during pre-mRNA splicing.,PTM:Extensively phosphorylated on serine residues in the RS domain.,similarity:Belongs to the splicing factor SR family.,similarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN., Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Cell Pathway	Nucleus speckle .
splicing.,PTM:Extensively phosphorylated on serine residues in the RS domain.,similarity:Belongs to the splicing factor SR family.,similarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2. Interacts with PNN., Background This gene encodes a member of the arginine/serine-rich splicing factor family.	Tissue Specificity	Aorta endothelial cell,Cerebrum,Epithelium,Lymph,
Background This gene encodes a member of the arginine/serine-rich splicing factor family. The encoded protein likely functions in mRNA processing. [provided by RefSeq,	Function	splicing.,PTM:Extensively phosphorylated on serine residues in the RS domain.,similarity:Belongs to the splicing factor SR family.,similarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Found in a pre-mRNA splicing complex with SFRS4, SFRS5, SNRNP70, SNRPA1, SRRM1 and SRRM2.
	Background	This gene encodes a member of the arginine/serine-rich splicing factor family. The encoded protein likely functions in mRNA processing. [provided by RefSeq,

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





Feb 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using SRp75 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd