



TEF Monoclonal Antibody

Catalog No	BYmab-02094
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	TEF
Protein Name	Thyrotroph embryonic factor
Immunogen	The antiserum was produced against synthesized peptide derived from human TEF. AA range:211-260
Specificity	TEF Monoclonal Antibody detects endogenous levels of TEF protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TEF; KIAA1655; Thyrotroph embryonic factor
Observed Band	45-55kD
Cell Pathway	Nucleus.
Tissue Specificity	Brain,Kidney,Retina,
Function	function:Transcription factor that binds to and transactivates the TSHB promoter. Binds to a minimal DNA-binding sequence 5'-[TC][AG][AG][TTA[TC][AG]-3'.,induction:Accumulates according to a robust circadian rhythm.,similarity:Belongs to the bZIP family.,similarity:Belongs to the bZIP family. PAR subfamily.,similarity:Contains 1 bZIP domain.,subunit:Binds DNA as a homodimer or a heterodimer. Can form a heterodimer with DBP.,
Background	TEF, PAR bZIP transcription factor(TEF) Homo sapiens This gene encodes a member of the PAR (proline and acidic amino acid-rich) subfamily of basic region/leucine zipper (bZIP) transcription factors. It is expressed in a broad range of cells and tissues in adult animals, however, during embryonic development,

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



TEF expression appears to be restricted to the developing anterior pituitary gland, coincident with the appearance of thyroid-stimulating hormone, beta (TSHB). Indeed, TEF can bind to, and transactivate the TSHB promoter. It shows homology (in the functional domains) with other members of the PAR-bZIP subfamily of transcription factors, which include albumin D box-binding protein (DBP), human hepatic leukemia factor (HLF) and chicken vitellogenin gene-binding protein (VBP); VBP is considered the chicken homologue of TEF. Different members of the subfamily can readily form heterodimers, and share DNA-binding, and transcriptional regulatory


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using TEF Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd