



TFE3 Monoclonal Antibody

Catalog No	BYmab-02101
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	TFE3
Protein Name	Transcription factor E3
Immunogen	The antiserum was produced against synthesized peptide derived from human TFE3. AA range:101-150
Specificity	TFE3 Monoclonal Antibody detects endogenous levels of TFE3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TFE3; BHLHE33; Transcription factor E3; Class E basic helix-loop-helix protein 33; bHLHe33
Observed Band	62kD
Cell Pathway	Cytoplasm, cytosol . Nucleus . When nutrients are present, phosphorylation by MTOR prevents nuclear translocation and activity (PubMed:22692423, PubMed:30733432). Conversely, inhibition of mTORC1, starvation and lysosomal disruption, promotes dephosphorylation and translocation to the nucleus (PubMed:22692423, PubMed:30733432). .
Tissue Specificity	Ubiquitous in fetal and adult tissues.
Function	disease:Chromosomal aberrations involving TFE3 are recurrent in alveolar soft part sarcoma (ASPS) [MIM:606243]. Translocation t(X;17)(p11;q25) with ASPSCR1 forms a ASPSCR1-TFE3 fusion protein.,disease:Chromosomal aberrations involving TFE3 are recurrent in alveolar soft part sarcoma (ASPS). Translocation t(X;17)(p11;q25) with ASPSCR1 forms a ASPSCR1-TFE3 fusion protein.,disease:Chromosomal aberrations involving TFE3 may be a cause of papillary renal cell carcinoma (PRCC) [MIM:605074]. Translocation

Nanjing BYabscience technology Co.,Ltd



t(X;1)(p11.2;q21.2) with PRCC; translocation t(X;1)(p11.2;p34) with PSF; inversion inv(X)(p11.2;q12) that fuses NONO to TFE3.,function:Positive-acting transcription factor that binds to the immunoglobulin enhancer MUE3 motif. It also binds very well to a USF/MLTF site. Binding of TFE3 to DNA induces DNA binding.,similarity:Belongs to the Mit/TFE family.,similarity:Contains 1 basic helix-loop-

Background

This gene encodes a basic helix-loop-helix domain-containing transcription factor that binds MUE3-type E-box sequences in the promoter of genes. The encoded protein promotes the expression of genes downstream of transforming growth factor beta (TGF-beta) signaling. This gene may be involved in chromosomal translocations in renal cell carcinomas and other cancers, resulting in the production of fusion proteins. Translocation partners include PRCC (papillary renal cell carcinoma), NONO (non-POU domain containing, octamer-binding), and ASPSCR1 (alveolar soft part sarcoma chromosome region, candidate 1), among other genes. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabs science technology Co.,Ltd