



TFEB Monoclonal Antibody

Catalog No	BYmab-02102
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	TFEB
Protein Name	Transcription factor EB
Immunogen	The antiserum was produced against synthesized peptide derived from human TFEB. AA range:10-59
Specificity	TFEB Monoclonal Antibody detects endogenous levels of TFEB protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TFEB; BHLHE35; Transcription factor EB; Class E basic helix-loop-helix protein 35; bHLHe35
Observed Band	52kD
Cell Pathway	Cytoplasm, cytosol . Lysosome membrane . Nucleus . Mainly present in the cytoplasm (PubMed:23434374, PubMed:33691586). Under aberrant lysosomal storage conditions, it translocates from the cytoplasm to the nucleus (PubMed:21617040, PubMed:22576015, PubMed:23434374). The translocation to the nucleus is regulated by ATP13A2 (PubMed:23434374, PubMed:27278822). Colocalizes with mTORC1 on the lysosomal membrane: when nutrients are present, phosphorylation by MTOR prevents nuclear translocation and activity (PubMed:22343943, PubMed:22692423). Conversely, inhibition of mTORC1, starvation and lysosomal disruption, promotes dephosphorylation and translocation to the nucleus (PubMed:22343943, PubMed:22692423). Exported from the nucleus in response to nutrient availability (PubMed:30120233). In macro
Tissue Specificity	B-cell,Brain,Kidney,Lung,Muscle,Spleen,

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com


官方热线: 025-5229-8998

监督电话: 15950492658



Function	function:Probable transcription factor binds to the USF/MLTF site and probably recognizes E-box sequences in the heavy-chain immunoglobulin enhancer.,similarity:Belongs to the MiT/TFE family.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,subunit:Efficient DNA binding requires dimerization with another bHLH protein.,
Background	function:Probable transcription factor binds to the USF/MLTF site and probably recognizes E-box sequences in the heavy-chain immunoglobulin enhancer.,similarity:Belongs to the MiT/TFE family.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,subunit:Efficient DNA binding requires dimerization with another bHLH protein.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using TFEB Monoclonal Antibody