



TFIIIA Monoclonal Antibody

Catalog No	BYmab-02108
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	GTF3A
Protein Name	Transcription factor IIIA
Immunogen	Synthesized peptide derived from the Internal region of human TFIIIA.
Specificity	TFIIIA Monoclonal Antibody detects endogenous levels of TFIIIA protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	GTF3A; Transcription factor IIIA; TFIIIA
Observed Band	41kD
Cell Pathway	Nucleus.
Tissue Specificity	Ubiquitous.
Function	caution:It is uncertain whether Met-1 is the initiator. Based on the lack of an in-frame AUG codon, mammalian TFIIIA may be translated from this non-AUG initiation site, which has a good Kozak context and which is well conserved among mammals.,function:Interacts with the internal control region (ICR) of approximately 50 bases within the 5S RNA genes, is required for correct transcription of these genes by RNA polymerase III. Also binds the transcribed 5S RNA's. May initiate transcription of the 5S ribosomal RNA gene and maintain the stability of transcription of other genes.,sequence caution:Translation N-terminally shortened.,similarity:Contains 9 C2H2-type zinc fingers.,tissue specificity:Ubiquitous.,

Nanjing BYabscience technology Co.,Ltd



Background

The product of this gene is a zinc finger protein with nine Cis[2]-His[2] zinc finger domains. It functions as an RNA polymerase III transcription factor to induce transcription of the 5S rRNA genes. The protein binds to a 50 bp internal promoter in the 5S genes called the internal control region (ICR), and nucleates formation of a stable preinitiation complex. This complex recruits the TFIIC and TFIIB transcription factors and RNA polymerase III to form the complete transcription complex. The protein is thought to be translated using a non-AUG translation initiation site in mammals based on sequence analysis, protein homology, and the size of the purified protein. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using TFIIIA Monoclonal Antibody