



Carbonyl Reductase 3 Monoclonal Antibody

Catalog No BYmab-02526 Isotype IgG Reactivity Human;Mouse;Rat Applications WB Gene Name CBR3 Protein Name Carbonyl reductase [NADPH] 3 Immunogen The antiserum was produced against synthesized peptide derived from human CBR3. AA range:151-200 Specificity Carbonyl Reductase 3 Monoclonal Antibody detects endogenous levels of Carbonyl Reductase 3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3 Observed Band 31kD Cell Pathway Cytoplasm . Tissue Specificity Dietected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function Catalytic activity:R-CHOH-R'		
Reactivity Human;Mouse;Rat Applications WB Gene Name CBR3 Protein Name Carbonyl reductase [NADPH] 3 Immunogen The antiserum was produced against synthesized peptide derived from human CBR3. AA range:151-200 Specificity Carbonyl Reductase 3 Monoclonal Antibody detects endogenous levels of Carbonyl Reductase 3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3 Observed Band 31kD Cell Pathway Cytoplasm. Tissue Specificity Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R¹ + NADP(+) = R-CO-R¹ + NADPH., similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family. Background Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [Provided by RefSeq,	Catalog No	BYmab-02526
Applications WB Gene Name CBR3 Protein Name Carbonyl reductase [NADPH] 3 Immunogen The antiserum was produced against synthesized peptide derived from human CBR3. AA range:151-200 Specificity Carbonyl Reductase 3 Monoclonal Antibody detects endogenous levels of Carbonyl Reductase 3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3 Observed Band 31kD Cell Pathway Cytoplasm. Tissue Specificity Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH, similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family., Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to annomeric NADPH-dependent is oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to annomeric NADPH-dependent is oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to annomeric NADPH-dependent is oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to annomeric NADPH-dependent is oxidoreductase.	Isotype	IgG
Gene Name CBR3 Protein Name Carbonyl reductase [NADPH] 3 Immunogen The antiserum was produced against synthesized peptide derived from human CBR3. AA range:151-200 Specificity Carbonyl Reductase 3 Monoclonal Antibody detects endogenous levels of Carbonyl Reductase 3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3 Observed Band 31kD Cell Pathway Cytoplasm . Tissue Specificity Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH., similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family. Background Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl c	Reactivity	Human;Mouse;Rat
Protein Name Carbonyl reductase [NADPH] 3 Immunogen The antiserum was produced against synthesized peptide derived from human CBR3. AA range:151-200 Specificity Carbonyl Reductase 3 Monoclonal Antibody detects endogenous levels of Carbonyl Reductase 3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3 Observed Band 31kD Cell Pathway Cytoplasm . Tissue Specificity Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH, similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family. Background Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding valored pounds of their correspondin	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human CBR3. AA range:151-200 Specificity Carbonyl Reductase 3 Monoclonal Antibody detects endogenous levels of Carbonyl Reductase 3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3 Observed Band 31kD Cell Pathway Cytoplasm . Tissue Specificity Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity: R-CHOH-R* + NADP(+) = R-CO-R* + NADPH., similarity: Belongs to the short-chain dehydrogenases/reductases (SDR) family Background Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilob	Gene Name	CBR3
CBR3. AA range:151-200 Specificity Carbonyl Reductase 3 Monoclonal Antibody detects endogenous levels of Carbonyl Reductase 3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3 Observed Band 31kD Cell Pathway Cytoplasm Tissue Specificity Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity: R-CHOH-R' + NADP(+) = R-CO-R' + NADPH., similarity: Belongs to the short-chain dehydrogenases/reductases (SDR) family., Background Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobasses and is closely linked to another carbonyl reductase or CBR1. [provided by RefSeq,	Protein Name	Carbonyl reductase [NADPH] 3
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3 Observed Band 31kD Cell Pathway Cytoplasm. Tissue Specificity Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family., Background CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq,	Immunogen	
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3 Observed Band 31kD Cell Pathway Cytoplasm . Tissue Specificity Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH., similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family., Background Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq,	Specificity	Carbonyl Reductase 3 Monoclonal Antibody detects endogenous levels of Carbonyl Reductase 3 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3 Observed Band 31kD Cell Pathway Cytoplasm Tissue Specificity Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH., similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family., Background Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq,	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3 Observed Band 31kD Cell Pathway Cytoplasm . Tissue Specificity Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family., Background Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductases. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq,	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3 Observed Band 31kD Cell Pathway Cytoplasm . Tissue Specificity Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family., Background Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq,	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3 Observed Band 31kD Cell Pathway Cytoplasm . Tissue Specificity Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family., Background Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq,	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3 Observed Band 31kD Cell Pathway Cytoplasm. Tissue Specificity Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH., similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family., Background Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq,	Concentration	1 mg/ml
Synonyms CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3 Observed Band 31kD Cell Pathway Cytoplasm. Tissue Specificity Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family., Background Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq,	Purity	≥90%
Observed Band Cell Pathway Cytoplasm . Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family., Background Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq,	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm . Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family., Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq,	Synonyms	CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3
Tissue Specificity Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family., Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq,	Observed Band	31kD
liver, spleen, leukocyte, prostate and testis. Function catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family., Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq,	Cell Pathway	Cytoplasm .
Background Carbonyl reductase 3 catalyzes the reduction of a large number of biologically and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq,	Tissue Specificity	Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis.
and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq,	Function	catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family.,
	Background	and pharmacologically active carbonyl compounds to their corresponding alcohols. The enzyme is classified as a monomeric NADPH-dependent oxidoreductase. CBR3 contains three exons spanning 11.2 kilobases and is closely linked to another carbonyl reductase gene - CBR1. [provided by RefSeq,

Nanjing BYabscience technology Co.,Ltd






matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Carbonyl Reductase 3 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd