



# COX11 Monoclonal Antibody

<b>Catalog No</b>	BYmab-02539
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	COX11
<b>Protein Name</b>	Cytochrome c oxidase assembly protein COX11 mitochondrial
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human COX11. AA range:51-100
<b>Specificity</b>	COX11 Monoclonal Antibody detects endogenous levels of COX11 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	COX11; Cytochrome c oxidase assembly protein COX11; mitochondrial
<b>Observed Band</b>	31kD
<b>Cell Pathway</b>	Mitochondrion inner membrane ; Single-pass membrane protein ; Intermembrane side .
<b>Tissue Specificity</b>	Ubiquitous.
<b>Function</b>	function:Exerts its effect at some terminal stage of cytochrome c oxidase synthesis, probably by being involved in the insertion of the copper B into subunit I.,similarity:Belongs to the COX11/ctaG family.,subunit:Interacts with CNNM4/ACDP4.,tissue specificity:Ubiquitous.,
<b>Background</b>	Cytochrome c oxidase (COX), the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may function in the regulation

**Nanjing BYabscience technology Co.,Ltd**



and assembly of the complex. This nuclear gene encodes a protein which is not a structural subunit, but may be a heme A biosynthetic enzyme involved in COX formation, according to the yeast mutant studies. However, the studies in *Rhodobacter sphaeroides* suggest that this gene is not required for heme A biosynthesis, but required for stable formation of the Cu(B) and magnesium centers of COX. This human protein is predicted to c


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using COX11 Monoclonal Antibody