



# CYP19A1 Monoclonal Antibody

Catalog No	BYmab-02562
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CYP19A1
Protein Name	Cytochrome P450 19A1
Immunogen	The antiserum was produced against synthesized peptide derived from human Cytochrome P450 19A1. AA range:221-270
Specificity	CYP19A1 Monoclonal Antibody detects endogenous levels of CYP19A1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CYP19A1; ARO1; CYAR; CYP19; Cytochrome P450 19A1; Aromatase; CYPXIX; Cytochrome P-450AROM; Estrogen synthase
Observed Band	53kD
Cell Pathway	Endoplasmic reticulum membrane ; Multi-pass membrane protein . Microsome membrane ; Multi-pass membrane protein .
Tissue Specificity	Widely expressed, including in adult and fetal brain, placenta, skin fibroblasts, adipose tissue and gonads.
Function	catalytic activity:RH + reduced flavoprotein + O(2) = ROH + oxidized flavoprotein + H(2)O.,cofactor:Heme group.,disease:Defects in CYP19A1 are a cause of familial gynecomastia [MIM:139300]. This is characterized by an estrogen excess due to an increased aromatase activity.,disease:Defects in CYP19A1 are the cause of aromatase deficiency (AROD) [MIM:107910]. AROD is a rare disease in which fetal androgens are not converted into estrogens due to placental aromatase deficiency. Thus, pregnant women exhibit a hirsutism, which spontaneously resolves after post-partum. At birth, female babies present with pseudohermaphroditism due to virilization of extern genital organs. In adult females, manifestations include delay of puberty, breast hypoplasia and primary

Nanjing BYabscience technology Co.,Ltd



amenorrhoea with multicystic ovaries.,function:Catalyzes the formation of aromatic C18 estrogens from C19 androgens.,online information:

#### Background

cytochrome P450 family 19 subfamily A member 1(CYP19A1) Homo sapiens  
This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and catalyzes the last steps of estrogen biosynthesis. Mutations in this gene can result in either increased or decreased aromatase activity; the associated phenotypes suggest that estrogen functions both as a sex steroid hormone and in growth or differentiation. Alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2014],


#### matters needing attention

Avoid repeated freezing and thawing!

#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

### Products Images



Western Blot analysis of various cells using CYP19A1 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd