




## CYP4X1 Monoclonal Antibody

Catalog No	BYmab-02604
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	CYP4X1
Protein Name	Cytochrome P450 4X1
Immunogen	The antiserum was produced against synthesized peptide derived from human Cytochrome P450 4X1. AA range:231-280
Specificity	CYP4X1 Monoclonal Antibody detects endogenous levels of CYP4X1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
-	
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year CYP4X1; Cytochrome P450 4X1; CYPIVX1
Storage Stability Synonyms Observed Band	-20°C/1 year  CYP4X1; Cytochrome P450 4X1; CYPIVX1  60kD  Endoplasmic reticulum membrane ; Single-pass membrane protein . Microsome
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year  CYP4X1; Cytochrome P450 4X1; CYPIVX1  60kD  Endoplasmic reticulum membrane; Single-pass membrane protein. Microsome membrane; Single-pass membrane protein.  Expressed in brain, heart, kidney and skin and, at lower levels, in skeletal muscle and liver (PubMed:16478468, PubMed:18549450). In the brain, high levels are detected in amygdala and lower levels in globus pallidus and cerebellum (PubMed:18549450). In the heart, very high levels in aorta, but very low levels in other heart regions (PubMed:16478468, PubMed:18549450). Also expressed in
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year  CYP4X1; Cytochrome P450 4X1; CYPIVX1  60kD  Endoplasmic reticulum membrane; Single-pass membrane protein. Microsome membrane; Single-pass membrane protein.  Expressed in brain, heart, kidney and skin and, at lower levels, in skeletal muscle and liver (PubMed:16478468, PubMed:18549450). In the brain, high levels are detected in amygdala and lower levels in globus pallidus and cerebellum (PubMed:18549450). In the heart, very high levels in aorta, but very low levels in other heart regions (PubMed:16478468, PubMed:18549450). Also expressed in breast, prostate and colon (PubMed:18549450).  catalytic activity:RH + reduced flavoprotein + O(2) = ROH + oxidized flavoprotein

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







steroids and other lipids. The expression pattern of a similar rat protein suggests that this protein may be involved in neurovascular function in the brain. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2016],

matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

