



DECR2 Monoclonal Antibody

Catalog No	BYmab-02613
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	DECR2
Protein Name	Peroxisomal 2,4-dienoyl-CoA reductase
Immunogen	The antiserum was produced against synthesized peptide derived from human DECR2. AA range:217-266
Specificity	DECR2 Monoclonal Antibody detects endogenous levels of DECR2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DECR2; PDCR; Peroxisomal 2; 4-dienoyl-CoA reductase; pDCR; 2,4-dienoyl-CoA reductase 2
Observed Band	36kD
Cell Pathway	Peroxisome .
Tissue Specificity	Brain,Epithelium,Lung,Testis,
Function	catalytic activity:Trans-2,3-didehydroacyl-CoA + NADP(+) = trans,trans-2,3,4,5-tetradehydroacyl-CoA + NADPH.,function:Auxiliary enzyme of beta-oxidation. Participates in the degradation of unsaturated fatty enoyl-CoA esters having double bonds in both even- and odd-numbered positions in peroxisome. Catalyzes the NADP-dependent reduction of 2,4-dienoyl-CoA to yield trans-3-enoyl-CoA. Has activity towards short and medium chain 2,4-dienoyl-CoAs, but also towards 2,4,7,10,13,16,19-docosaheptaenoyl-CoA, suggesting that it does not constitute a rate limiting step in the peroxisomal degradation of docosahexaenoic acid.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family.,similarity:Belongs to the short-chain

Nanjing BYabscience technology Co.,Ltd



dehydrogenases/reductases (SDR) family. 2,4-dienoyl-CoA reductase subfamily.,

Background

catalytic activity:Trans-2,3-didehydroacyl-CoA + NADP(+) = trans,trans-2,3,4,5-tetradehydroacyl-CoA + NADPH.,function:Auxiliary enzyme of beta-oxidation. Participates in the degradation of unsaturated fatty enoyl-CoA esters having double bonds in both even- and odd-numbered positions in peroxisome. Catalyzes the NADP-dependent reduction of 2,4-dienoyl-CoA to yield trans-3-enoyl-CoA. Has activity towards short and medium chain 2,4-dienoyl-CoAs, but also towards 2,4,7,10,13,16,19-docosaheptaenoyl-CoA, suggesting that it does not constitute a rate limiting step in the peroxisomal degradation of docosahexaenoic acid.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family. 2,4-dienoyl-CoA reductase subfamily.,


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd