



FX Monoclonal Antibody

Catalog No BYmab-02637 Isotype IgG Reactivity Human;Mouse;Rat Applications WB Gene Name TSTA3 Protein Name GDP-L-fucose synthase Immunogen The antiserum was produced against synthesized peptide derived from hard and a range:221-270	uman
Reactivity Human;Mouse;Rat Applications WB Gene Name TSTA3 Protein Name GDP-L-fucose synthase Immunogen The antiserum was produced against synthesized peptide derived from h TSTA3. AA range:221-270	uman
Applications WB Gene Name TSTA3 Protein Name GDP-L-fucose synthase Immunogen The antiserum was produced against synthesized peptide derived from h TSTA3. AA range:221-270	uman
Gene Name TSTA3 Protein Name GDP-L-fucose synthase Immunogen The antiserum was produced against synthesized peptide derived from h TSTA3. AA range:221-270	uman
Protein Name GDP-L-fucose synthase Immunogen The antiserum was produced against synthesized peptide derived from h TSTA3. AA range:221-270	uman
Immunogen The antiserum was produced against synthesized peptide derived from h TSTA3. AA range:221-270	uman
TSTA3. AA range:221-270	uman
EVAN I IA (1) I I I I I I I I I I I I I I I I I I I	
Specificity FX Monoclonal Antibody detects endogenous levels of FX protein.	
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azio	de.
Source Monoclonal, Mouse, IgG	
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.	
Dilution WB 1:500-2000	
Concentration 1 mg/ml	
Purity ≥90%	
Storage Stability -20°C/1 year	
Synonyms TSTA3; SDR4E1; GDP-L-fucose synthase; GDP-4-keto-6-deoxy-D-mann 5-epimerase-4-reductase; Protein FX; Red cell NADP(H)-binding protein; Short-chain dehydrogenase/reductase family 4E member 1	
Observed Band 40kD	
Cell Pathway cytoplasm,cytosol,extracellular exosome,	
Tissue Specificity Erythrocyte,Lung,Placenta,Skin,	
catalytic activity:GDP-L-fucose + NADP(+) = GDP-4-dehydro-6-deoxy-D-mannose + NADPH.,function:Two step NADP-dependent conversion of GDP-4-dehydro-6-deoxy-D-mannose to GDP-fucose, involving an epimerase and a reductase reaction.,pathway:Nucleotide-sugar biosynthesis; GDP-L-fucose biosynth de novo pathway; GDP-L-fucose from GDP-D-mannose: step 2/2.,similarity:Belongs to the fucose synthetase family.,subunit:Homodime	

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background	Tissue specific transplantation antigen P35B is a NADP(H)-binding protein. It catalyze the two-step epimerase and the reductase reactions in GDP-D-mannose metabolism, converting GDP-4-keto-6-D-deoxymannose to GDP-L-fucose. GDP-L-fucose is the substrate of several fucosyltransferases involved in the expression of many glycoconjugates, including blood group ABH antigens and developmental adhesion antigens. Mutations in this gene may cause leukocyte adhesion deficiency, type II. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658