



LPAAT-ε Monoclonal Antibody

Catalog No	BYmab-02660
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	AGPAT5
Protein Name	1-acyl-sn-glycerol-3-phosphate acyltransferase epsilon
Immunogen	The antiserum was produced against synthesized peptide derived from human AGPAT5. AA range:241-290
Specificity	LPAAT- ε Monoclonal Antibody detects endogenous levels of LPAAT- ε protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	AGPAT5; 1-acyl-sn-glycerol-3-phosphate acyltransferase epsilon; 1-acylglycerol-3-phosphate O-acyltransferase 5; 1-AGP acyltransferase 5; 1-AGPAT 5; Lysophosphatidic acid acyltransferase epsilon; LPAAT-epsilon
Observed Band	45kD
Cell Pathway	Endoplasmic reticulum membrane ; Multi-pass membrane protein . Nucleus envelope . Mitochondrion .
Tissue Specificity	Widely expressed.
Function	catalytic activity:Acyl-CoA + 1-acyl-sn-glycerol 3-phosphate = CoA + 1,2-diacyl-sn-glycerol 3-phosphate.,caution:It is uncertain whether Met-1 or Met-12 is the initiator.,domain:The HXXXX motif is essential for acyltransferase activity and may constitute the binding site for the phosphate moiety of the glycerol-3-phosphate.,function:Converts lysophosphatidic acid (LPA) into phosphatidic acid by incorporating an acyl moiety at the sn-2 position of the glycerol backbone.,pathway:Phospholipid metabolism; CDP-diacylglycerol biosynthesis; CDP-diacylglycerol from sn-glycerol 3-phosphate: step 2/3.,similarity:Belongs to the 1-acyl-sn-glycerol-3-phosphate acyltransferase

Nanjing BYabscience technology Co.,Ltd



family.,

Background

This gene encodes a member of the 1-acylglycerol-3-phosphate O-acyltransferase family. This integral membrane protein converts lysophosphatidic acid to phosphatidic acid, the second step in de novo phospholipid biosynthesis. A pseudogene of this gene is present on the Y chromosome. [provided by RefSeq, Aug 2014],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabs science technology Co.,Ltd