



## Malic enzyme 1 Monoclonal Antibody

Catalog No         BYmab-02665           Isotype         IgG           Reactivity         Human;Mouse;Rat           Applications         WB           Gene Name         ME1           Protein Name         NADP-dependent malic enzyme           Immunogen         The antiserum was produced against synthesized peptide derived from human ME1. AA range:468-517           Specificity         Malic enzyme 1 Monoclonal Antibody detects endogenous levels of Malic enzyme 1 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         ME1; NADP-dependent malic enzyme; NADP-ME; Malic enzyme 1           Observed Band         64kD           Cell Pathway         Cytoplasm.           Tissue Specificity         Expressed in all tissues tested including liver, placenta and white adipose tissue.           Function         catalytic activity; (S)-malate + NADP(+) = pyruvate + CO(2) + NADPH, cofactor: Divalent metal cations. Pref		
Reactivity         Human;Mouse;Rat           Applications         WB           Gene Name         ME1           Protein Name         NADP-dependent malic enzyme           Immunogen         The antiserum was produced against synthesized peptide derived from human ME1. AA range:468-517           Specificity         Malic enzyme 1 Monoclonal Antibody detects endogenous levels of Malic enzyme 1 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         ME1; NADP-dependent malic enzyme; NADP-ME; Malic enzyme 1           Observed Band         64kD           Cell Pathway         Cytoplasm .           Tissue Specificity         Expressed in all tissues tested including liver, placenta and white adipose tissue.           Function         catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH, cofactor: Divalent metal cations. Prefers magnesium or manganese, online information: The Singapore human mutation and polymorphism database, similarity:Belongs to the malic enzy	Catalog No	BYmab-02665
Applications  Gene Name  ME1  Protein Name  NADP-dependent malic enzyme  Immunogen  The antiserum was produced against synthesized peptide derived from human ME1. AA range:468-517  Specificity  Malic enzyme 1 Monoclonal Antibody detects endogenous levels of Malic enzyme 1 protein.  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  ME1; NADP-dependent malic enzyme; NADP-ME; Malic enzyme 1  Observed Band  64kD  Cell Pathway  Cytoplasm.  Tissue Specificity  Expressed in all tissues tested including liver, placenta and white adipose tissue.  Function  catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH, cofactor: Divalent metal cations. Prefers magnesium or manganese, online information: The Singapore human mutation and polymorphism database, similarity. Pelongs to the malic enzymes family, subunit: Homotetramer., tissue specificity: Expressed in all tissues tested including liver, placenta and white adipose tissue.  Background  This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxy/alton of malate, links the divolovitic and citric acid cycles. The	Isotype	IgG
Gene Name ME1 Protein Name NADP-dependent malic enzyme  Immunogen The antiserum was produced against synthesized peptide derived from human ME1. AA range:468-517 Specificity Malic enzyme 1 Monoclonal Antibody detects endogenous levels of Malic enzyme 1 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms ME1; NADP-dependent malic enzyme; NADP-ME; Malic enzyme 1  Observed Band 64kD  Cell Pathway Cytoplasm .  Tissue Specificity Expressed in all tissues tested including liver, placenta and white adipose tissue.  Function Catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH, cofactor:Divalent metal cations. Prefers magnesium or manganese, online information:The Singapore human mutation and polymorphism database, similarity. Belongs to the malic enzymes family, subunit-Homotetramer, tissue specificity/Expressed in all tissues tested including liver, placenta and white adipose tissue.  Background This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the divocivitic acid cycles. The	Reactivity	Human;Mouse;Rat
Protein Name  NADP-dependent malic enzyme  The antiserum was produced against synthesized peptide derived from human ME1. AA range:468-517  Specificity  Malic enzyme 1 Monoclonal Antibody detects endogenous levels of Malic enzyme 1 protein.  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  ME1; NADP-dependent malic enzyme; NADP-ME; Malic enzyme 1  Observed Band  64kD  Cell Pathway  Cytoplasm  Tissue Specificity  Expressed in all tissues tested including liver, placenta and white adipose tissue.  Function  catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH, cofactor:Divalent metal cations. Prefers magnesium or manganese, online information:The Singapore human mutation and polymorphism database, similarity:Bellongs to the malic enzymes family, subunit-Homotetramer, tissue specificity:Expressed in all tissues tested including liver, placenta and white adipose tissue.  Background  This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxvlation of malate, links the glycolytic and citric acid cycles. The	Applications	WB
Immunogen         The antiserum was produced against synthesized peptide derived from human ME1. AA range:468-517           Specificity         Malic enzyme 1 Monoclonal Antibody detects endogenous levels of Malic enzyme 1 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         ME1; NADP-dependent malic enzyme; NADP-ME; Malic enzyme 1           Observed Band         64kD           Cell Pathway         Cytoplasm .           Tissue Specificity         Expressed in all tissues tested including liver, placenta and white adipose tissue.           Function         catalytic activity: (S)-malate + NADP(+) = pyruvate + CO(2) + NADP-H., cofactor: Divalent metal cations. Prefers magnesium or manganese, online information: The Singapore human mutation and polymorphism database, similarity: Belongs to the malic enzymes family, subunit: Homotetramer, tissue specificity: Expressed in all tissues tested including liver, placenta and white adipose tissue.           Background         This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH to refatty acid biosynthesis	Gene Name	ME1
ME1. AA range:468-517  Specificity Malic enzyme 1 Monoclonal Antibody detects endogenous levels of Malic enzyme 1 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms ME1; NADP-dependent malic enzyme; NADP-ME; Malic enzyme 1  Observed Band 64kD  Cell Pathway Cytoplasm .  Tissue Specificity Expressed in all tissues tested including liver, placenta and white adipose tissue.  Function catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH, cofactor:Divalent metal cations. Prefers magnesium or manganese., online information: The Singapore human mutation and polymorphism database, similarity:Belongs to the malic enzymes family, subunit:Homotetramer, tissue specificity:Expressed in all tissues tested including liver, placenta and white adipose tissue.  Background This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the divotic and critic acid cycles. The	Protein Name	NADP-dependent malic enzyme
Tissue Specificity  Expressed in all tissues tested including liver, placenta and white adipose tissue.  Source  Another Specificity  Expressed in all tissues tested including liver, placenta and white adipose tissue.  Background  This gene encodes a cytosolic, NADP-dependent enzyme that generates NADP-dependent enzyme, the reversible oxidative decarboxylation of malate, links the divolvic and citric acid cycles. The	Immunogen	
Source       Monoclonal, Mouse, IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       ME1; NADP-dependent malic enzyme; NADP-ME; Malic enzyme 1         Observed Band       64kD         Cell Pathway       Cytoplasm .         Tissue Specificity       Expressed in all tissues tested including liver, placenta and white adipose tissue.         Function       catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH, cofactor:Divalent metal cations. Prefers magnesium or manganese, online information:The Singapore human mutation and polymorphism database, similarity:Belongs to the malic enzymes family, subunit:Homotetramer, tissue specificity:Expressed in all tissues tested including liver, placenta and white adipose tissue.         Background       This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the divcolvtic and citric acid cycles. The	Specificity	Malic enzyme 1 Monoclonal Antibody detects endogenous levels of Malic enzyme 1 protein.
Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  ME1; NADP-dependent malic enzyme; NADP-ME; Malic enzyme 1  Observed Band  64kD  Cell Pathway  Cytoplasm.  Tissue Specificity  Expressed in all tissues tested including liver, placenta and white adipose tissue.  Function  Catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH.,cofactor:Divalent metal cations. Prefers magnesium or manganese.,online information: The Singapore human mutation and polymorphism database, similarity:Belongs to the malic enzymes family.,subunit:Homotetramer.,tissue specificity:Expressed in all tissues tested including liver, placenta and white adipose tissue.  Background  This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the dycolytic and citric acid cycles. The	Formulation	•
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms ME1; NADP-dependent malic enzyme; NADP-ME; Malic enzyme 1  Observed Band 64kD  Cell Pathway Cytoplasm .  Tissue Specificity Expressed in all tissues tested including liver, placenta and white adipose tissue.  Function catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH.,cofactor:Divalent metal cations. Prefers magnesium or manganese, online information:The Singapore human mutation and polymorphism database, similarity:Belongs to the malic enzymes family, subunit:Homotetramer., tissue specificity:Expressed in all tissues tested including liver, placenta and white adipose tissue.  Background This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the glycolytic and citric acid cycles. The	Source	Monoclonal, Mouse,lgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       ME1; NADP-dependent malic enzyme; NADP-ME; Malic enzyme 1         Observed Band       64kD         Cell Pathway       Cytoplasm .         Tissue Specificity       Expressed in all tissues tested including liver, placenta and white adipose tissue.         Function       catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH., cofactor:Divalent metal cations. Prefers magnesium or manganese.,online information:The Singapore human mutation and polymorphism database,similarity:Belongs to the malic enzymes family., subunit:Homotetramer., tissue specificity:Expressed in all tissues tested including liver, placenta and white adipose tissue.         Background       This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the glycolytic and citric acid cycles. The	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms ME1; NADP-dependent malic enzyme; NADP-ME; Malic enzyme 1  Observed Band 64kD  Cell Pathway Cytoplasm .  Tissue Specificity Expressed in all tissues tested including liver, placenta and white adipose tissue.  Function catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH.,cofactor:Divalent metal cations. Prefers magnesium or manganese.,online information:The Singapore human mutation and polymorphism database,similarity:Belongs to the malic enzymes family.,subunit:Homotetramer.,tissue specificity:Expressed in all tissues tested including liver, placenta and white adipose tissue.  Background This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the glycolytic and citric acid cycles. The	Dilution	WB 1:500-2000
Storage Stability  -20°C/1 year  ME1; NADP-dependent malic enzyme; NADP-ME; Malic enzyme 1  Observed Band  64kD  Cell Pathway  Cytoplasm.  Expressed in all tissues tested including liver, placenta and white adipose tissue.  Function  catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH.,cofactor:Divalent metal cations. Prefers magnesium or manganese.,online information:The Singapore human mutation and polymorphism database,similarity:Belongs to the malic enzymes family.,subunit:Homotetramer.,tissue specificity:Expressed in all tissues tested including liver, placenta and white adipose tissue.,  Background  This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the glycolytic and citric acid cycles. The	Concentration	1 mg/ml
Synonyms ME1; NADP-dependent malic enzyme; NADP-ME; Malic enzyme 1  Observed Band 64kD  Cell Pathway Cytoplasm .  Tissue Specificity Expressed in all tissues tested including liver, placenta and white adipose tissue.  Function catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH.,cofactor:Divalent metal cations. Prefers magnesium or manganese.,online information: The Singapore human mutation and polymorphism database,similarity:Belongs to the malic enzymes family.,subunit:Homotetramer.,tissue specificity:Expressed in all tissues tested including liver, placenta and white adipose tissue.,  Background This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the glycolytic and citric acid cycles. The	Purity	≥90%
Observed Band  Cell Pathway  Cytoplasm .  Tissue Specificity  Expressed in all tissues tested including liver, placenta and white adipose tissue.  Function  catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH.,cofactor:Divalent metal cations. Prefers magnesium or manganese.,online information:The Singapore human mutation and polymorphism database,similarity:Belongs to the malic enzymes family.,subunit:Homotetramer.,tissue specificity:Expressed in all tissues tested including liver, placenta and white adipose tissue.,  Background  This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the glycolytic and citric acid cycles. The	Storage Stability	-20°C/1 year
Cell Pathway  Cytoplasm .  Expressed in all tissues tested including liver, placenta and white adipose tissue.  Function  catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH.,cofactor:Divalent metal cations. Prefers magnesium or manganese.,online information:The Singapore human mutation and polymorphism database,similarity:Belongs to the malic enzymes family.,subunit:Homotetramer.,tissue specificity:Expressed in all tissues tested including liver, placenta and white adipose tissue.,  Background  This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the glycolytic and citric acid cycles. The	Synonyms	ME1; NADP-dependent malic enzyme; NADP-ME; Malic enzyme 1
Tissue Specificity  Expressed in all tissues tested including liver, placenta and white adipose tissue.  Catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH.,cofactor:Divalent metal cations. Prefers magnesium or manganese.,online information:The Singapore human mutation and polymorphism database,similarity:Belongs to the malic enzymes family.,subunit:Homotetramer.,tissue specificity:Expressed in all tissues tested including liver, placenta and white adipose tissue.,  Background  This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the glycolytic and citric acid cycles. The	Observed Band	64kD
Function  catalytic activity:(S)-malate + NADP(+) = pyruvate + CO(2) + NADPH.,cofactor:Divalent metal cations. Prefers magnesium or manganese.,online information:The Singapore human mutation and polymorphism database,similarity:Belongs to the malic enzymes family.,subunit:Homotetramer.,tissue specificity:Expressed in all tissues tested including liver, placenta and white adipose tissue.,  This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the glycolytic and citric acid cycles. The	Cell Pathway	Cytoplasm .
NADPH.,cofactor:Divalent metal cations. Prefers magnesium or manganese.,online information:The Singapore human mutation and polymorphism database,similarity:Belongs to the malic enzymes family.,subunit:Homotetramer.,tissue specificity:Expressed in all tissues tested including liver, placenta and white adipose tissue.,  Background  This gene encodes a cytosolic, NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the glycolytic and citric acid cycles. The	Tissue Specificity	Expressed in all tissues tested including liver, placenta and white adipose tissue.
NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the glycolytic and citric acid cycles. The	Function	NADPH.,cofactor:Divalent metal cations. Prefers magnesium or manganese.,online information:The Singapore human mutation and polymorphism database,similarity:Belongs to the malic enzymes family.,subunit:Homotetramer.,tissue specificity:Expressed in all tissues tested
	Background	NADPH for fatty acid biosynthesis. The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the glycolytic and citric acid cycles. The

Nanjing BYabscience technology Co.,Ltd


网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





	from elevated levels of thyroid hormones or by higher proportions of carbohydrates in the diet. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Nanjing BYabscience technology Co.,Ltd