



## **USP45** Monoclonal Antibody

Catalog No         BYmab-02840           Isotype         IgG           Reactivity         Human;Mouse;Rat           Applications         WB           Gene Name         USP45           Protein Name         Ubiquitin carboxyl-terminal hydrolase 45           Immunogen         The antiserum was produced against synthesized peptide derived from human USP45. AA range:301-350           Specificity         USP45 Monoclonal Antibody detects endogenous levels of USP45 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         USP45; Ubiquitin carboxyl-terminal hydrolase 45; Deubiquitinating enzyme 45; Ubiquitin thioesterase 45; Ubiquitin-specific-processing protease 45           Observed Band         90kD           Cell Pathway         Photoreceptor inner segment . Cytoplasm . Nucleus .           Tissue Specificity         Widely expressed . High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic ne		
Reactivity  Applications  WB  Gene Name  USP45  Protein Name  Ubiquitin carboxyl-terminal hydrolase 45  Immunogen  The antiserum was produced against synthesized peptide derived from human USP45. AA range:301-350  Specificity  USP45 Monoclonal Antibody detects endogenous levels of USP45 protein.  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  290%  Storage Stability  -20°C/1 year  Synonyms  USP45; Ubiquitin carboxyl-terminal hydrolase 45; Deubiquitinating enzyme 45; Ubiquitin thioesterase 45; Ubiquitin-specific-processing protease 45  Observed Band  90kD  Cell Pathway  Photoreceptor inner segment . Cytoplasm . Nucleus .  Tissue Specificity  Widely expressed . High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low expressed at high levels in the optic nerve, sclera and retina, with relatively low expressed at high levels in the optic nerve, sclera and retina, with relatively low expressed (lens and retinal pigment epithelium (PubMed:30573563).  Function  Catalytic activity: Ubiquitin C-terminal thioester + H(2)0 = ubiquitin + a thiol. similarity:Belongs to the peptidase C19 family, similarity:Contains 1 UBP-type zinc finger, tissue specificity:Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.  Background	Catalog No	BYmab-02840
Applications  Gene Name  USP45  Protein Name  Ubiquitin carboxyl-terminal hydrolase 45  Immunogen  The antiserum was produced against synthesized peptide derived from human USP45. AA range:301-350  Specificity  USP45 Monoclonal Antibody detects endogenous levels of USP45 protein.  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  290%  Storage Stability  -20°C/1 year  Synonyms  USP45; Ubiquitin carboxyl-terminal hydrolase 45; Deubiquitinating enzyme 45; Ubiquitin thioesterase 45; Ubiquitin-specific-processing protease 45  Observed Band  90kD  Cell Pathway  Photoreceptor inner segment . Cytoplasm . Nucleus .  Tissue Specificity  Widely expressed . High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function  Catalytic activity. Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol. similarity. Belongs to the peptidase C19 family, similarity. Contains 1 UBP-type zinc finger, tissue specificity. Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.  The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a critical requilator of DNA repair processes. and the	Isotype	IgG
Protein Name USP45  Protein Name Ubiquitin carboxyl-terminal hydrolase 45  Immunogen The antiserum was produced against synthesized peptide derived from human USP45. AA range:301-350  Specificity USP45 Monoclonal Antibody detects endogenous levels of USP45 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity 290%  Storage Stability -20°C/1 year  Synonyms USP45; Ubiquitin carboxyl-terminal hydrolase 45; Deubiquitinating enzyme 45; Ubiquitin thioesterase 45; Ubiquitin-specific-processing protease 45  Observed Band 90kD  Cell Pathway Photoreceptor inner segment . Cytoplasm . Nucleus .  Tissue Specificity Widely expressed. High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol. similarity Belongs to the peptidase C19 family. similarity. Contains 1 UBP-type zinc finger, tissue specificity. Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.  The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a rotical requilator of DNA repair endonuclease.	Reactivity	Human;Mouse;Rat
Protein Name Ubiquitin carboxyl-terminal hydrolase 45  Immunogen The antiserum was produced against synthesized peptide derived from human USP45. AA range:301-350  Specificity USP45 Monoclonal Antibody detects endogenous levels of USP45 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms USP45; Ubiquitin carboxyl-terminal hydrolase 45; Deubiquitinating enzyme 45; Ubiquitin thioesterase 45; Ubiquitin-specific-processing protease 45  Observed Band 90kD  Cell Pathway Photoreceptor inner segment . Cytoplasm . Nucleus .  Tissue Specificity Widely expressed. High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function catalytic activity: Ubiquitin C-terminal thioester + H(2)0 = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family, similarity: Contains 1 UBP-type zinc finger, tissue specificity: Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.  The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair processes, and the	Applications	WB
Immunogen  The antiserum was produced against synthesized peptide derived from human USP45. AA range:301-350  Specificity  USP45 Monoclonal Antibody detects endogenous levels of USP45 protein.  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source  Monoclonal, Mouse,lgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  USP45; Ubiquitin carboxyl-terminal hydrolase 45; Deubiquitinating enzyme 45; Ubiquitin thioesterase 45; Ubiquitin-specific-processing protease 45  Observed Band  90kD  Cell Pathway  Photoreceptor inner segment . Cytoplasm . Nucleus .  Tissue Specificity  Widely expressed . High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function  Videly expressed to the peptidase C19 family, similarity:Delongs to the peptidase C19 family, similarity:Contains 1 UBP-type zinc finger, tissue specificity:Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.  The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair processes, and the	Gene Name	USP45
USP45. AA range:301-350  Specificity USP45 Monoclonal Antibody detects endogenous levels of USP45 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity 290%  Storage Stability -20°C/1 year  Synonyms USP45; Ubiquitin carboxyl-terminal hydrolase 45; Deubiquitinating enzyme 45; Ubiquitin thioesterase 45; Ubiquitin-specific-processing protease 45  Observed Band 90kD  Cell Pathway Photoreceptor inner segment . Cytoplasm . Nucleus .  Tissue Specificity Widely expressed . High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, solera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function Catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol. similarity: Belongs to the peptidase C19 family., similarity: Contains 1 UBP-type zinc finger, tissue specificity: Broadly expressed, with highest levels in ovary, skeletal muscle and spleen., The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a critical regulator of DNA repair processes, and the	Protein Name	Ubiquitin carboxyl-terminal hydrolase 45
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms USP45; Ubiquitin carboxyl-terminal hydrolase 45; Deubiquitinating enzyme 45; Ubiquitin thioesterase 45; Ubiquitin-specific-processing protease 45  Observed Band 90kD  Cell Pathway Photoreceptor inner segment . Cytoplasm . Nucleus .  Tissue Specificity Widely expressed. High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity:Belongs to the peptidase C19 family., similarity:Contains 1 UBP-type zinc finger, tissue specificity:Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.  The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a critical regulator of DNA repair processes, and the	Immunogen	
Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms USP45; Ubiquitin carboxyl-terminal hydrolase 45; Deubiquitinating enzyme 45; Ubiquitin thioesterase 45; Ubiquitin-specific-processing protease 45  Observed Band 90kD  Cell Pathway Photoreceptor inner segment . Cytoplasm . Nucleus .  Tissue Specificity Widely expressed . High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiolsimilarity: Belongs to the peptidase C19 familysimilarity: Contains 1 UBP-type zinc fingertissue specificity: Broadly expressed, with highest levels in ovary, skeletal muscle and spleen  Background The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a critical regulator of DNA repair endonuclease.	Specificity	USP45 Monoclonal Antibody detects endogenous levels of USP45 protein.
Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  USP45; Ubiquitin carboxyl-terminal hydrolase 45; Deubiquitinating enzyme 45; Ubiquitin thioesterase 45; Ubiquitin-specific-processing protease 45  Observed Band  90kD  Cell Pathway  Photoreceptor inner segment . Cytoplasm . Nucleus .  Tissue Specificity  Widely expressed. High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function  catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity: Belongs to the peptidase C19 family., similarity: Contains 1 UBP-type zinc finger.,tissue specificity:Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.  Background  The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a critical regulator of DNA repair processes, and the	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms USP45; Ubiquitin carboxyl-terminal hydrolase 45; Deubiquitinating enzyme 45; Ubiquitin thioesterase 45; Ubiquitin-specific-processing protease 45  Observed Band 90kD  Cell Pathway Photoreceptor inner segment . Cytoplasm . Nucleus .  Tissue Specificity Widely expressed. High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,similarity:Contains 1 UBP-type zinc finger.,tissue specificity:Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.,  Background The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair processes, and the	Source	Monoclonal, Mouse,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       USP45; Ubiquitin carboxyl-terminal hydrolase 45; Deubiquitinating enzyme 45; Ubiquitin-specific-processing protease 45         Observed Band       90kD         Cell Pathway       Photoreceptor inner segment . Cytoplasm . Nucleus .         Tissue Specificity       Widely expressed. High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).         Function       catalytic activity; Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity:Belongs to the peptidase C19 family., similarity:Contains 1 UBP-type zinc finger., tissue specificity:Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.,         Background       The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a critical regulator of DNA repair processes, and the	Purification	·
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms USP45; Ubiquitin carboxyl-terminal hydrolase 45; Deubiquitinating enzyme 45; Ubiquitin thioesterase 45; Ubiquitin-specific-processing protease 45  Observed Band 90kD  Cell Pathway Photoreceptor inner segment . Cytoplasm . Nucleus .  Tissue Specificity Widely expressed. High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 familly., similarity: Contains 1 UBP-type zinc finger., tissue specificity: Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.,  The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a critical regulator of DNA repair processes, and the	Dilution	WB 1:500-2000
Storage Stability  -20°C/1 year  USP45; Ubiquitin carboxyl-terminal hydrolase 45; Deubiquitinating enzyme 45; Ubiquitin thioesterase 45; Ubiquitin-specific-processing protease 45  Observed Band  90kD  Cell Pathway  Photoreceptor inner segment . Cytoplasm . Nucleus .  Widely expressed . High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function  catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., similarity: Contains 1 UBP-type zinc finger., tissue specificity: Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.,  The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a critical regulator of DNA repair processes, and the	Concentration	1 mg/ml
Synonyms  USP45; Ubiquitin carboxyl-terminal hydrolase 45; Deubiquitinating enzyme 45; Ubiquitin thioesterase 45; Ubiquitin-specific-processing protease 45  Observed Band  90kD  Cell Pathway  Photoreceptor inner segment . Cytoplasm . Nucleus .  Widely expressed. High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function  catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., similarity: Contains 1 UBP-type zinc finger., tissue specificity: Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.,  The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a critical regulator of DNA repair processes, and the	Purity	≥90%
Ubiquitin thioesterase 45; Ubiquitin-specific-processing protease 45  90kD  Cell Pathway  Photoreceptor inner segment . Cytoplasm . Nucleus .  Widely expressed. High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function  catalytic activity: Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol., similarity: Belongs to the peptidase C19 family., similarity: Contains 1 UBP-type zinc finger., tissue specificity: Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.,  The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a critical regulator of DNA repair processes, and the	Storage Stability	-20°C/1 year
Cell Pathway  Photoreceptor inner segment . Cytoplasm . Nucleus .  Widely expressed. High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function  catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,similarity:Contains 1 UBP-type zinc finger.,tissue specificity:Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.,  The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a critical regulator of DNA repair processes, and the	Synonyms	
Tissue Specificity  Widely expressed. High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function  catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,similarity:Contains 1 UBP-type zinc finger.,tissue specificity:Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.,  The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a critical regulator of DNA repair processes, and the	Observed Band	90kD
expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).  Function  catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,similarity:Contains 1 UBP-type zinc finger.,tissue specificity:Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.,  The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a critical regulator of DNA repair processes, and the	Cell Pathway	Photoreceptor inner segment . Cytoplasm . Nucleus .
thiol., similarity: Belongs to the peptidase C19 family, similarity: Contains 1 UBP-type zinc finger, tissue specificity: Broadly expressed, with highest levels in ovary, skeletal muscle and spleen.,  The protein encoded by this gene is a deubiquitylase that binds ERCC1, the catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a critical regulator of DNA repair processes, and the	Tissue Specificity	Widely expressed. High expression is detected in the cerebellum. In the eye, it is expressed at high levels in the optic nerve, sclera and retina, with relatively low levels in the choroid, lens and retinal pigment epithelium (PubMed:30573563).
catalytic subunit of the XPF-ERCC1 DNA repair endonuclease. This endonuclease is a critical regulator of DNA repair processes, and the	Function	thiol.,similarity:Belongs to the peptidase C19 family, similarity:Contains 1 UBP-type zinc finger.,tissue specificity:Broadly expressed, with highest levels in
		evaly, excitat massic and opioent,

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



repair ability of XPF-ERCC1. [provided by RefSeq, Sep 2016],


matters needing		
attention		

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using USP45 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd